

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

62.19

LIBRARY
RECEIVED

MAY 19 1945 ★

U S Department of Agriculture


EVERETT'S

Garden Annual - 1945

EVERETT SEED COMPANY, ATLANTA, GA.

78 Alabama St., S.W. (Between Whitehall and Pryor)

...with a
**COMPLETE
GARDENING
GUIDE**




SEE INSIDE
FRONT COVER
for descriptions
of items illus-
trated here

A very different kind of seed catalog!

As you'll soon see, this Garden Guide is far more than the usual seed catalog. It lists the necessary seeds and supplies, but it goes a lot farther than that . . . by giving you many pages of good, practical garden information . . . just the kind of facts every home gardener seeks. Then, too, this book doesn't come to you simply because you are a "name" on the list of some far-distant seed house. It is sent by a seedsman in your own home territory . . . to ask—


"What can we do to help you, neighbor?"

You see, we've been right here in this section for a long time. We know the soil . . . the climate. Our experience qualifies us to help you as no far-away seed house can. Besides, we're here the year round, ready to serve you and work with you . . . as all good neighbors do.



BROCCOLI, Green Sprouting

This fine vegetable is extremely rich in vitamins, and should be in every garden. Tender heads, delicious flavor.




TOMATO, Marglobe

A second-early, red-fruited tomato. Medium size fruits, smooth, solid and deep, relatively free from cracks. Resistant to wilt and rust.




BEAN, Tendergreen

A distinctly superior stringless, green-podded bush bean. Pods are rich dark green, meaty, smooth, very tender and finely flavored. Strong, vigorous, early.



LETTUCE, Iceberg

Very desirable for home gardens. Sure heading. Makes compact heads, with white interior. Crisp and sweet. Resistant to heat. Late variety.



SQUASH, Early White Bush

Fruits greenish when young, become white at maturity. 3-in. long, 7½ to 8½-in. wide. 53 days. A real treat for your table.

FRONT COVER FLOWERS and VEGETABLES

COSMOS, Sensation Dazzler—An early flowering strain that grows about 4 ft. high. Has giant blooms, 4 to 6 inches across, in a lovely velvety crimson shade. Wavy, fluted petals. Larger flowers may be obtained by disbudding.

ONION, Prizetaker—This mild, light straw colored onion of the Spanish type becomes very large under good growing conditions, sometimes weighing up to 4 lbs. Very solid, crisp and juicy with a thin skin.

ZINNIA, Giants of California, Mixed—Beautiful, large double flowers, 5 to 6 in. across. They are smooth topped with the petals lying flat upon each other. Vigorous, well-branched plants with many long, strong flower stems. Ideal for cutting.

POPPY, California—Easily cultured, dainty flowers well adapted for beds and edgings. Blooms early and continues until frost. The modern strains come in many delicate tints and have lovely petals of satiny texture.

TOMATO, Jubilee—A bright, orange colored fruit, with few seeds and very high in Vitamins A and C. Fine for slicing because of uniformity and solidity. Won bronze medal, 1943.

ASTER, Giant Crego Navy Blue—Deep, purplish blue flowers with long, fully double, curled petals on 20-in. branching plants. The blooms are very large and attractive. A popular new strain.

CORN, Golden Bantam—Undeniably the favorite for table use because of its tenderness and sweetness. The cob is small but unsurpassed for quality. Matures in about 80 days.

SPRING SUGGESTION NO. 1—Get it all down on paper . . . here and now!

This year, don't take chances with your shopping memory! Go through this book carefully, and jot down everything you'll need for your spring planting. We've left these two pages blank just to help you do this. After your list is finished, drop in, and give it to us. We'll either fill your order while you wait . . . or have it ready when you wish.

Make your 1945 garden shopping easier, more pleasant, more complete! Get all your seeds, supplies and equipment needs down on paper, right now! Hand your list to us . . . and we'll do the rest.

Everett Seed Co.

GARDEN SHOPPING LIST

All gardeners of experience know that success with bulbs, etc., depends largely on weather conditions and proper cultivation, so in accordance with the universal custom of the seed trade we give no warranty, express or implied, as to the productiveness

of any seeds, bulbs or plants we sell and will not be in any way responsible for the crop. Our liability, in all instances, is limited to the purchase price of the seed, bulbs or plants.

FARM SEEDS? Let Everett supply you!

Everett's Farm, Field and Grass Seed Department is ready to take care of you, this year as always. Our stocks are complete—and we're looking forward to your calls.

For over thirty years this experienced, conscientious seed house has been supplying the seed varieties and strains best adapted to this section. We can't give you, here, the complete list and description of the many items we stock. But we can assure you that our list is very complete.

Just ask for any Farm, Field or Grass Seed varieties you don't find on these pages. We can take care of all your needs.

We particularly invite you to visit this store—and discuss your plans and your needs with us. We are always more than glad to talk things over—and to give you our careful advice.

Sudan
Millet
Field Corn
Alfalfa
Hegari
Kaffir
Milo
Crotalaria
Austrian Peas
Vetch
Lupine
Oats
Rye
Barley
Wheat

Cow Peas Soy Beans Peanuts

Velvet Beans Crowder Peas Lespedeza

Clover Pasture Grasses Sorghum Cane

LAWN GRASS SEEDS



*A better lawn
for your home?*

WE CAN HELP YOU!

For years it has been our business to provide exactly the right kind of seed for each particular lawn condition.

We know lawns—and we know this territory! So bring any lawn troubles you may have to us. Our long experience has taught us answers to the particular problems of this locality—and we're glad to pass them on to you.

Remember, too—you can depend on us for the greatest bargain in lawn seed—the most grass of the best type for each dollar you spend!

HOW MUCH SEED?

Many lawn failures are due to the common mistake of using too little seed!

It takes plenty of seed to get the thick, lasting turf you want—and to crowd out weeds.

On new lawns, you need 5 pounds of seed per 1,000 square feet. Don't risk your lawn success by stinting on seed! The cost is so small it isn't worth it.

No lawn mixture can meet the needs of every territory—or all the varying requirements in the same vicinity.

To be sure of having a thick, lasting turf, you must sow a mixture of true turf grasses properly suited to your own soil and climatic conditions. To get such a mixture, buy your seed from a seedsman right here in your own territory.

Best Lawn Mixture

Our selected lawn mixture is the very finest combination of high quality seeds . . . a marvelous mixture that years of experimenting has proved to be safe and sound. It produces a lawn that is like a carpet of green velvet . . . smooth, even, and thick.

Bermuda Grass Unhulled—As valuable for lawns or golf courses as it is for pasturage—or wherever a thick, heavy sod is desired. Needs plenty of water. When Bermuda grass turns brown in late fall or winter, give it a good application of lawn fertilizer.

Hulled Bermuda Grass—The highest grade of Bermuda, with the hard, almost water-tight hulls removed. Produces much quicker and more certain results. Germination, with proper moisture conditions, in 1 to 2 weeks.


Carpet Grass—Exceedingly hardy. Does better in poor, sandy soil than any other grass. The more mowing and tramping it gets, the better it likes it.

Winter Rye—The grass that withstands winter weather perfectly. Plant it in fall for a nice green lawn during the winter months. An annual—must be planted every year.

Red Top—Particularly suitable for wet, low spots, as it stands wet weather, or even overflowing, better than other grasses. Also does well on almost any good, rich soil. Often thrives where blue grass fails.

For shady spots

FINEST SHADE MIXTURE—Wherever shade makes it difficult to obtain a good lawn, our special shade mixture is heartily recommended. Excellent quality, shade-enduring grasses and clovers . . . blended into a mixture that gives surprising results.



Our 1945 Lawn Seed Prices

BERMUDA GRASS: 1 lb. 90c; 5 lbs. \$4.25. Prices fluctuate. Ask for quantity prices.

RED TOP: 1 lb. 35c; 5 lbs. \$1.50; 10 lbs. \$2.90; 25 lbs. \$7.00; 100 lbs. \$27.00.

CARPET GRASS: 1 lb. 55c; 5 lbs. \$2.50; 10 lbs. \$4.50; 25 lbs. \$11.00; 100 lbs. \$42.00.

HULLED BERMUDA GRASS: 1 lb. \$1.45; 5 lbs. \$7.00. Ask for prices on larger quantities.

RYE GRASS: 1 lb. 20c; 5 lbs. 85c; 10 lbs. \$1.50; 25 lbs. \$3.50; 100 lbs. \$12.50.

KENTUCKY BLUE GRASS: 1 lb. 50c; 5 lbs. \$2.00; 10 lbs. \$3.80; 25 lbs. \$9.25; 100 lbs. \$36.00.

DRUID HILLS EVERGREEN LAWN MIXTURE: 1 lb. 40c; 5 lbs. \$1.90; 10 lbs. \$3.70; 25 lbs. \$9.00; 100 lbs. \$35.00.

SHADY LAWN MIXTURE: 1 lb. 45c; 5 lbs. \$2.15; 10 lbs. \$4.20; 25 lbs. \$10.25; 100 lbs. \$40.00.

**EVERETT SEED CO.
ATLANTA, GEORGIA**

EVERETT'S 1945 Vegetable Seed Prices

All prices subject to change without notice; all stocks offered subject to previous sale.

		Doz.	100	
		Lb.	5 Lbs. per Lb.	
ASPARAGUS Roots, 2-year....		\$0.40	\$2.75	
BEANS, Green Podded, Dwarf Bush—All varieties	\$0.40	\$0.75	\$1.50	
Red Valentine Bush Bean.....	.40	.75	1.50	
BEANS, Wax Podded, Dwarf Bush—All varieties40	.75	1.50	
BEANS, Pole—All varieties....	.40	.75	1.50	
BEANS, Lima, Dwarf—All varieties40	.75	1.50	
BEANS, Lima, Pole: All varieties40	.75	1.50	
King of Garden (large).....	.40	.75	1.50	
Oz. 4 Oz. 8 Oz. Lb.				
BEETS—Detroit Dark Red.....	\$0.20	\$0.65	\$1.20	\$2.25
All other varieties.....	.20	.60	1.10	2.00
SWISS CHARD—Lucullus15	.40	.75	1.25
MANGEL WURZEL15	.50	.90	1.75
BROCCOLI35	1.00	1.85	3.75
BRUSSELS SPROUTS			(10c packets only)	
CABBAGE—Chinese25	.75	1.35	2.50
Copenhagen Market40	1.25	2.25	4.00
All other varieties listed.....	.35	1.10	2.00	3.50
CARROTS—All varieties listed.....	.20	.60	1.10	2.00
CAULIFLOWER			(25c packets only)	
CELERY			(10c packets only)	
SWEET CORN—Country Gentleman	½ Lb. Lb. 2 Lbs. Per Lb.	\$0.25 \$0.45	\$0.75	
Stowells Evergreen25	.45	.75	
Golden Cross Bantam.....	.30	.50	.90	
Golden Bantam25	.45	.75	
Cherokee Sweet25	.45	.75	
Truckers Favorite10	.20	.35	
POPCORN—Yellow Dynamite..	Oz. 4 Oz. 8 Oz. Lb.	.20 .35	.50	
COLLARD—All varieties	\$0.10	\$0.25	\$0.45	\$0.75
CUCUMBER: A. & C.....	.25	.75	1.40	2.50
Improved Long Green.....	.25	.75	1.40	2.50
All other varieties.....	.20	.60	1.10	2.00
DILL—Long Island Mammoth..			(10c packets only)	
EGGPLANT—All varieties.....	½ Oz. Oz. 4 Oz. Lb.	\$0.35 \$0.60	\$2.00	\$7.00
ENDIVE—All varieties15	.25	.65	2.00
KALE—All varieties20	.60	2.00
LETTUCE—Great Lakes (New)		.50	1.75	6.00
Imperial varieties30	.90	3.25
New York30	.90	3.25
Iceberg20	.60	2.00
All others listed.....		.20	.60	2.00
COS OR ROMAINE.....		.25	.75	
MUSKMELON—Honey Dew....		.25	.75	2.25
All other varieties.....		.20	.60	1.90
WATERMELON—Dixie Queen.		.20	.50	1.50
Florida Giant or Cannon Ball.		.25	.65	1.90
Stone Mountain20	.50	1.50
All other varieties.....		.15	.40	1.25
MUSTARD—Tendergreen Mustard Spinach15	.40	1.25

In Packets at 10c per Packet

except as otherwise noted


All other varieties.....	.10	.30	.95
OKRA—All varieties10	.30	.95
PARSLEY20	.50	1.40
PARSNIPS25	.75	2.25
PEAS—All varieties	Lb. 2 Lbs. Per Lb.	\$0.40 \$0.75	\$1.50
PEPPERS—California Wonder.	½ Oz. Oz. 4 Oz.	\$0.30 \$0.55	\$1.90
Large Bell or Bull Nose.....	.25	.45	1.50
Pimento20	.30	.90
Long Red Cayenne—Hot.....	.25	.45	1.50
(Unable to supply other varieties, due to crop failure)			
Oz. 4 Oz. 8 Oz. Lb.			
PUMPKIN—Kentucky Field or Large Cheese	\$0.15	\$0.30	\$0.55
All other varieties.....	.20	.45	.85
RADISH—All varieties10	.30	.50
SPINACH—New Zealand.....	.25	.85	
All other varieties.....	.10	.25	.40
SQUASH—All varieties15	.45	.80
TOMATO—Brimmer75	2.50	4.50
Ponderosa75	2.50	4.50
Oxheart			(10c packets only)
Improved Stone45	1.20	2.00
Marglobe Wilt resistant.....	.45	1.20	2.00
Pritchard45	1.20	2.00
Stokesdale45	1.20	2.00
Rutgers45	1.20	2.00
All other varieties.....	.60	1.75	3.25
TURNIP—Seven Top10	.25	.40
All other varieties.....	.10	.25	.40
RUTABAGA or SWEDE TURNIP10	.25	.40

Grow your own vitamins!

This nation has the finest food supply in the world. Yet nutrition experts have proved that most of us are poorly nourished. As we all know now, even our "three good squares a day" often do not give us certain minerals and vitamins that we need . . . now more than ever.

These necessary elements may be bought at the drug store. But they can also be home grown, at far less cost! Garden fresh vegetables can provide nearly all of them, in ample quantity.

That's why the offerings on this page are the world's finest bargain in vitamins.



Selected VEGETABLE SEEDS

ASPARAGUS

Mary Washington—The best of several rust resistant strains recently developed. Large, firm shoots. Green with purple tops. Very fine flavor.

BEANS

Green Podded, Dwarf or Bush

Black Valentine Stringless—Hardy, early maturing, very productive. Pods, dark green, 6-6½ in. long, brittle, smooth, stringless. Seeds black. 51 days.

Bountiful—An early, heavy-yielding variety. The leading flat podded bean for market and home use. Flat, light-green, stringless, but slightly fibrous pods of finest quality. 49 days.

Giant Stringless Green Pod—Large, sturdy plants. Meaty, round pods, brittle and strictly stringless at all stages. Medium green pods, oval, yellowish-brown seed. 54 days.

Stringless Green Pod—Early, sturdy. Very meaty, round, fully stringless and fibreless. 50 days.

Tendergreen—Large, erect plants. Abundant, round, dark-green pods, strictly stringless and fleshy. Brown seeds blotched with light fawn. 54 days.

Wax Podded, Dwarf or Bush Varieties

Golden Wax—Small, erect plants, moderately productive. Straight, thick, flat pods, creamy yellow and stringless. 51 days.

All Vegetable Seeds
10c PER PACKET
UNLESS OTHERWISE NOTED

In bulk quantities at money-saving prices.
Special quotations to market gardeners.
Prices subject to change without notice.

Pencil Pod Black Wax—Early and productive. Rich yellow, deep curved round pods. 5½ to 6-in. long. Stringless, tender, brittle and fleshy. 52 days.

Round Pod Kidney Wax or Brittle Wax—Large, erect, vigorous plants. Medium yellow curved pods, crease-backed, entirely stringless, very brittle, fleshy and succulent. 58 days.


Stringless Kidney Wax (Oval Pod)—Excellent mid-season variety, suitable especially for home and market garden use. 5½ to 7 in. long. 55 days.

Sure Crop—Also called Bountiful Wax. Golden yellow pods 6 in. to 7 in. long. Stringless, fine quality. 55 days.

Beans, Pole

Genuine Cornfield (Striped Creaseback)—Vigorous-growing, prolific. Pods round, straight, light green; fleshy, tender, but somewhat stringy. 71 days.

ESPECIALLY RECOMMENDED



COLLARD Georgia or Southern

Favorite of winter "greens." Grows to 2 ft., forming large, loose heads of succulent leaves on long stem. 24 to 36 days.

Ideal Market or Black Valentine Pole—Earliest pole bean. Plant hardy, good climber. Pods round, silvery green, brittle, tender, stringless when young. Black seed. 60 days.

Kentucky Wonder—Strong climber, hardy, long-bearing. Curved, almost round pods. Slightly stringy, but brittle and fibreless. Meaty. 65 days.

McCaslan—Dry shelled or snap, good climber. Medium green thick, flat pods. Meaty, stringless and tender when young. 65 days.

Stringless Black Creaseback.

Lima, Dwarf or Bush

Baby Potato—1940 All-America. Small, thick-seeded butter bean with real flavor, early maturity and prolific growth. 12 to 16-in. tall.

Fordhook Bush—Straight pods with plump, large beans of excellent quality. 75 days.

About our selection of vegetable seeds

You'll find a hand-picked list of vegetable seed varieties offered on these pages. They are the kinds we know from experience are the most suited to home gardening.

Of course, you'll find many longer lists than we've given you here. But out of the many thousands of varieties available, we've simply chosen the best of the lot for quality, yield, and your own gardening needs.

If there should be some varieties you want, and can't find listed in this book . . . ask us about them.

We can either supply what you want . . . or get it for you.

Bountiful

Tendergreen

Pencil Pod
Black Wax

Giant
Stringless


Improved
Golden Wax

Kentucky
Wonder

**TRUE-TO-SCALE
ILLUSTRATIONS**

showing correct
comparison of
size and form


Early Jersey Wakefield




Copenhagen Market



Chinese Cabbage,
Chihili



Golden Acre



CHINESE
CABBAGE
Chihili



PEAS, Improved
Telephone



EGGPLANT
Black Beauty




SQUASH
Fordhook
Zucchini



SQUASH
Zucchini Bush



CANTALOUPE, Hearts of Gold



Give your garden a head start! with big, healthy VEGETABLE PLANTS

This year, treat your family to earlier vegetables by starting your garden with our Specially Selected Plants. These are big, healthy, strong plants that will grow and thrive, and produce abundantly the vitamin-rich, healthful vegetables everyone craves and needs.

We can supply you with most every type of thriving vegetable plant, ready to set out. They are all husky specimens . . . hardy and full of growth. This year, have earlier and more vegetables by setting out PLANTS.

BEANS

Lima, Dwarf or Bush (Cont'd)

Henderson's Bush—Known as Baby Lima or Butter Bean. Plants small, early and bushy. 65 days. Good for home gardener; very productive.

Jackson Wonder (Speckled Bush)—Pods medium size with 4 to 5 thick, large beans. Drought resistant and very prolific. Dry beans mottled in color. 66 days.

Lima, Pole

Florida Butter Bean—Dependable, popular, prolific. 78 days.

Sieva, Carolina or Small White—Plant 10 to 12-ft. tall, and bears over a long season. Medium green pods with 3 to 4 beans of excellent quality. 77 days.

BEETS

Crosby's Egyptian—Widely grown for early beets. Flattened globe shaped roots with small tap root. Excellent quality, tender and sweet. 60 days.

Detroit Dark Red—Standard of excellence in table beets. Smooth, globular roots of deep ox-blood red—sweet and tender. 68 days.

Early Blood Turnip—Medium early. Tops medium, somewhat coarse. Bright red. 65 to 70 days.

Early Wonder—Early variety. Semi-globular tender, blood-red flesh. 58 days.

Stock Beets or Mangel Wurzel—See Mangel Wurzel.

BROCCOLI

Italian Green Sprouting—Plant bears a succession of sprouting heads about 5 in. long, which, if kept cut, will be replaced by others for 8 to 10 weeks. 55 to 65 days.

BRUSSELS SPROUTS

Dwarf Improved—Firm, thick-set tender heads. Easily grown. Early, dependable.

CABBAGE

Standard Early Varieties

Charleston or Large Wakefield—Heart-shaped heads nearly a week later than Jersey Wakefield and half

again as large. 7 to 8-in. long. 70 to 75 days.

Copenhagen Market—Excellent, early short season type. Short stems, 6½-in., 3½ to 4-lb. heads. Solid, round and of good quality with few outer leaves, stems short. 65 to 70 days.

Golden Acre—Extra early Copenhagen type. Dwarf plants, small round solid, 6-in., 3-lb. heads. Globular in shape and with attractive smooth leaves. May be set close together. 63 to 65 days.

Jersey Wakefield—Early, with small conical heads, 7-in. long, short stems, 5-in., 2 to 2½-lb. heads. 64 days.

Intermediate Varieties

Early Flat Dutch—Good second early variety. Plants medium. Very vigorous. Large, flat solid heads 6-in. deep, 11 to 12-in. in diameter, 11 to 12-lb. 90 days.

Glory of Enkhuizen—Attractive, round solid heads of finest quality. Weigh 6 to 8 lbs. Finely ribbed, tender leaves. Excellent for kraut, slaw, etc. 80 days.

Marion Market—Large, firm, round heads, used for early kraut. A development from Copenhagen Market. 7-in., 4-lb. heads. 75 to 80 days.

Late or Winter Varieties

Ferry's Hollander—Medium sized round heads. Firm and solid. Wonderful quality. Best variety for home gardeners to plant for winter storage. 100 days.

Savoy—Large, flat heads with dark green, crinkled leaves. Flavor is mild and appetizing. Weigh 6 to 7 lbs. 90 days.

Red Varieties

Mammoth Red Rock—Best of the red cabbages. Hard, round purplish-red heads. Good keeper, 6 to 8-in., 7 to 8-lb. heads. 100 days.

CHINESE CABBAGE

Chihili or Improved Pekin—Dependable, early heading. The most popular variety. Solid tapered heads, 3 to 4-in. thick, 18 to 20-in. long. 75 days.

CARROTS

Chantenay, Red Cored—Root has thick shoulder and tapers to slight stump root, 5-in. long. Red cored. 70 days.

Danvers Half Long—Sweet, tender roots, 6 to 7-in. long tapering to a blunt point. Fine for storing. 75 days.

Imperator—Fine-grained, tender. Rich orange, indistinct core. Roots smooth, tapered to semi-blunt. 77 days.

Long Orange—Heavy cropper. Roots taper to a point. Red-orange flesh, core slightly lighter. 88 days.

CAULIFLOWER

Snowball—Favorite early variety. Pure white compact heads of fine flavor, 6 to 7-in. 55 days. Pkt. 25c.

CELERY

Golden Self Blanching—Compact plants with yellowish-green foliage. Eroad solid stalks, nutty flavor. Blanches readily. 120 days.

COLLARD

Cabbage Collard—Combines qualities of cabbage and collard, but finer flavored than either.

Southern or Georgia—An excellent cooked green. Long stemmed plant, with clustered leaves. Withstands heat and bad soil. 24 to 36 days.

SWEET CORN

Country Gentleman or Shoe Peg—Prolific late variety. Favored by canners. Irregular kernels, very deep, slender, sweet. 93 days.

Golden Bantam—The best and most favorably known of all the yellow varieties. Stalks often have 2 ears. Ears 8 rowed, kernels broad, with tender hull, sweet and of very fine flavor. 80 days.

Golden Cross Hybrid—Extremely uniform. Ears 10 to 14 rows, slightly lighter yellow than Golden Bantam. Highly resistant to Stewart's disease. 85 days.

Ioana Hybrid—1940 All-America. Ears 7½ to 8 in., well filled with 12 or 14 rows of deep, medium narrow, light yellow kernels. Highly resistant to drought and bacterial wilt. 87 days.

Stowell's Evergreen—White. Large ears about 8-in. long; big kernels, very sweet. One of the best late varieties. 93 days.

Tendergold—This hybrid was developed as a high quality second-early yellow hybrid for canners. Ears are cylindrical to slightly tapering, 7-8 in. long. Golden Bantam type. 80 days.

All Vegetable Seeds


10c PER PACKET

UNLESS OTHERWISE NOTED

In bulk quantities at money-saving prices.
Special quotations to market gardeners.
Prices subject to change without notice.



SQUASH, Early
Prolific Straightneck



CARROT
Nantes



CORN
Ioana

GARDEN OR ROASTING EAR CORN

Adams Improved or Large Early—Large-eared roasting variety, earlier than Trucker's Favorite, 12 to 14 rowed. Kernels white and tender when young. 70 to 73 days.

Trucker's Favorite—Extensively used for roasting ears and shipping. Ears 12 to 16 rowed. Kernels white, fairly tender, and moderately sweet. 70 to 75 days. Plant early or late.

POP CORN

Yellow Dynamite—Rich, creamy yellow, smooth and round kernels, which pop double size of others. Ears 8 to 9 in. long. Vigorous grower. 115 days.

CUCUMBER

A and C—A long dark green cucumber which merits a trial. Uniform, nearly cylindrical, well rounded at ends.

Cubit—Won Bronze Medal Award in 1944 All-American Selections. Dark green, white spine, cylindrical with blunt ends. Firm flesh and uniform coloring. New and worthwhile. 25c pkt.

Improved Long Green—Black spined. Weighs about $3\frac{3}{4}$ lbs. Best for slicing or pickling. Very hardy and disease resistant. 62 days.

Pickling—Largest of the pickling varieties, also good for slicing. 59 days.

Straight Eight, w.s.—Unsurpassed as slicing cucumber. All-American Selection. $2\frac{1}{2}$ -lb. fruits rounded at ends, medium green. 68 days.

DILL

Long Island Mammoth—Used for flavoring. Flat seeds have strong, bitter flavor.

EGGPLANT

Black Beauty—Large, egg-shaped, smooth, very dark purple fruits. Keep well. Plant bears 4 to 5 fruits. Early. 80 days.

Florida Highbush—Vigorous plant, tall and upright growth. Resists drought, blight and wet weather better than most other varieties. Cylindrical in shape. 85 days.

New Hampshire Hybrid—All-America Silver Medal 1939. The dark purple fruits are of highest quality.

ENDIVE

Deep Heart Fringed—All-America Silver Medal, 1940. An entirely new type of endive. Leaf margins deeply cut and curled. Instead of lying flat on

the ground, outer leaves turn upward, making this variety less susceptible to bottom rot than most others. Well-fitted hearts, closely packed with abundance of fine leaves, blanch to clear creamy yellow. Suitable for home and market garden. 70 days.

Green Curled Ruffec—Plants 16 to 18 in. in diameter. Fine fall variety. White, tender, fleshy. 95 days.

KALE or BORECOLE

Dwarf Green Curled—Hardy, large and very attractive bright, deep green. 100 days.

Early Curled Siberian (Sprouts)—A vigorous growing variety of low spreading habit. Grayish green leaves which are only slightly curled at edges. Extremely hardy.

KOHLRABI

Early Vienna—8 to 10-in. leaves on slender stems. Bulbs 2 to 3-in., globular, light green. Crisp, tender, clear white flesh. 55 to 60 days.

LETTUCE

Heading or Cabbage

Big Boston—Popular for cold frame forcing and outside culture. Medium, compact heads with creamy yellow heart. Smooth, glossy leaves, edges wavy, and slightly tinged with reddish brown. 75 days.

Great Lakes — Bronze Medal, 1944, A.A.S. Iceberg type lettuce newly developed. Large, solid heads resistant to heat and sun. 25c per pkt.

Imperial No. 44 (Improved Iceberg)

—A sure leader even during dry hot weather. Heads medium large, slightly flattened, extremely compact. 82 days.

Imperial No. 847 (Florida Iceberg)—Heads are of good size and solid. Does well in hot weather and is resistant to tip burn. Very hardy. 83 days.

New York—Large heads, well blanched, sweet, tender. 80 days.

Loose Leaved Varieties

Black Seeded Simpson—Light green, frilled and crumpled. 45 days.

Chicken—Produces largest amount of leaves of any lettuce. Fast-growing. For poultry and rabbit feeding. 40 days.

Simpson's Early Curled—Hardy and dependable. Non-heading. Leaves large frilled, crumpled, light green; form a compact bunch at center. 45 days.

Cos or Romaine

White Paris or Trianon—Medium large self-folding, dark green loaf shaped heads. Greenish-white, well blanched interior. 66 days.

MANGEL WURZEL

Mammoth Long Red—Very popular, 30 to 50 tons per acre. Roots grow half above the ground. Light red, flesh white with rose tinge. 110 days.

MELONS

Muskmelon and Cantaloupe

Banana—Fruits smooth and slender, 20 inches long and 4 inches thick, resembling a banana. Flesh salmon color, with banana-like flavor. 90 days.

Hale's Best, No. 36—Improved. Early, uniform, oval, light ribbed, heavily netted. Thick, salmon flesh. Sweet, spicy. 80 days.

Hearts of Gold—Nearly round fruit, weight 2 pounds. Distinctly ribbed, deep green with fine grey netting. Thick flesh, deep salmon in color, tender and sweet. 94 days.

Especially RECOMMENDED


PEPPER, World Beater

A large, handsome pepper that grows to 5 in. long, $3\frac{1}{2}$ in. diameter. Three- or four-lobed, with very thick, mild and sweet flesh. Fruit is glossy green changing to dull red when ripe. Plants are vigorous and productive.





ENDIVE, Deep Heart Fringed

Favorite of French chefs for green salads. This improved strain has full, deep, compact heart and thick, green leaves curled at the edges, blanched at the center. Can be sown early, as soon as ground can be worked, or later for main crop.



THIS YEAR
it's FLOWERS, too!
Vegetables? Of course! They grow dollars in return for pennies! They're your duty! But in a world longing for peace, grow flowers, too. The world needs beauty . . . now more than ever!





Muskmelon (Continued)

Honey Ball—Small round fruit, weight 3½ lbs. Yellowish-white, smooth, fine netting. Flesh is grey-green in color, very sweet and juicy. 104 days.

Honey Dew—The well-known green-fleshed melon for warm climates. Smooth, ivory colored skin. Flesh is sweet and juicy. 112 days.

Improved Perfecto—Roundish fruit, weight 2½ lbs. No ribs. Completely covered with hard grey netting. Flesh is thick and salmon colored, sweet and juicy. 90 days.

Rocky Ford, Golden Lined—Fruits small, with rather large seed cavity. Nearly round, weight 2½ lbs. No ribs. Heavily covered with hard grey netting. Flesh thick, green with gold tinge at center. Very juicy and spicy. Excellent quality. 92 days.

Watermelon

Dixie Queen—Very prolific. Bright red, crisp, splendid quality fibreless flesh. Very few small, white seeds. 85 days.

Florida Favorite—Oblong shaped. Dark green rind with light green stripes. Bright red and very sweet. 85 days.

Florida Giant—A huge melon, growing sometimes to weight of 100 lbs. Oval-round with blunt ends, it has dark green skin and thick, tough rind. Flesh is of excellent quality, firm and red. 90 days.

Leesburg Wilt Resistant—A recently developed strain of Kleckley's Sweet. Long, dark green melon. Very uniform and excellent shipping variety. 90 days.

Stone Mountain—Also called Dixie Belle. High quality shipping variety. Fruits very large, oval-round, blunt ends. Dark green, tough rind. Flesh rich scarlet, fine grained, sweet. Seed white with black tips. 90 days.

Tom Watson—Red heart strain. Large fruit, uniform cylindrical, with tough, elastic rind, faintly veined. 90 days.

Wondermelon or Kleckley's Sweet—Large, cylindrical, slightly ribbed fruits. Deep red, juicy, sweet flesh. 85 days.

MUSTARD

Florida Broad Leaf—Vigorous, upright plants. Leaves large, broad and deep green; quick grower. 43 days.

Chinese Broadleaf—Large, oblong, slightly crumpled leaves, scalloped at edges. Medium green, vigorous plants. 45 days.

Mustard Spinach or Tendergreen—Wholesome, quick-growing green, combining mustard and spinach flavors. 28 days.

Southern Giant Curled Longstanding—Large, frilled and crumpled. Light green leaves. 35 days.

OKRA or GUMBO

Clemson Spineless—All-America Silver Medal Award for 1939. Pods are rich green, straight, uniform, ridged and of best quality. 56 days.

Dwarf Green—Early, prolific dwarf growing sort. Dark green fluted, 5 to 7-in. pointed pods. Most desirable, compact growing okra. 50 days.

Perkins' Mammoth Long Green—Early and prolific. Very tender, long pods. Deep green and slightly corrugated. 68 days.

White Velvet—Early and prolific. Pods pale greenish-white, 6 to 7-in. long. Meaty, tender. Strong, tall-growing, 3½ feet in height. 62 days.

ONIONS

Yellow Varieties

Prizetaker—Large, globe shaped, with glossy, thin straw-colored skin. Coarse, mild flesh. Sometimes weigh as much as 4 lbs. each. 102 to 104 days.

Sweet Spanish—Large, globe-shaped, exceptionally mild and sweet—excellent for eating raw or boiled. Light yellow skin and pure white, firm, crisp, and tasty flesh. 112 days.

Yellow Bermuda—Very early, medium sized, flat. Thin loose skin. Flesh nearly white, mild and sweet. The most widely grown Bermuda Onion. 92-94 days.

Red Varieties

Red Creole—Grown largely in the South. Large, solid semi-globe of reddish hue and fine grain. Unexcelled for home garden. Good keeper.

Red Globe—Finest of all red onions. Perfectly round, good sized bulbs with thick, small necks. Skin is deep purplish-red. Strong flavored flesh, white tinged with pink. 112-114 days.

White Varieties

Crystal Wax Bermuda—Sometimes called White Bermuda. Bulbs flat, medium-sized, clear white. 92 to 94 days.

White Portugal or Silverskin—Excellent flat variety. Dependable cropper, excellent for sets, for green bunching, or as a pickler. 100 to 102 days.

ONION SETS

The quickest way to get onions early. We offer choice, select, dry sets, of highest quality. (Prices on application.)

WHITE RED YELLOW

PARSNIP

Hollow Crown—The most generally grown kind. Roots 2½ to 3-in. thick at shoulder. 12 to 14-in. long, uniformly tapered, hollow crowned. 95 days.

PARSLEY

Moss Curled or Triple Curled—Compact, dark green leaves, curled and fine-cut. 70 days.

PEAS

Early and Second Variety

Alaska—Wilt resistant. Used for canning and early home garden. Blunt, straight, light-green, 3-in. pods. Plants 32-in. 60 days.

Laxtonian—Exceptionally choice and prolific. Dark green, rather coarse vines, producing straight pointed, well-filled 2½-in. pods. 62 days.

Laxton's Progress—Medium dark green vine, 16 to 18-in. Single pods almost an inch wide and 4½ to 5-in. long, with 7 to 9 large peas. 62 days.

Little Marvel—Outstanding dwarf. Fine quality, large yield. Single and double, dark green, 3-in. pods—blunt, plump, well filled with 7 to 8 medium sized, light green tender peas. 62 days.

Thomas Laxton—Medium height. Strong vine bears numerous dark green, blunt ended pods, 4 to 4½-in. One of earliest. 57 days.

White Marrowfat—Vines grow 5 feet tall. Pods plump, blunt, straight, light green, about 3-in. long. 81 days.

Peas — Later Varieties

Everbearing—Long season, wilt resistant, marrowy peas. Pods plump, blunt, straight. 32-in. vine. 76 days.

Tall Telephone, Dark Podded—Mid-season favorite, high quality. 74 days.

For earlier onions...better onions...plant SETS



Grow Onions from Sets

For the home gardener, growing onions from sets saves labor, makes good results more certain, gives you onions ready to use (as green onions) in 2 to 3 weeks and mature onions about a month earlier than those grown from seed.

Plant as soon as ground can be worked. Set in 12-in. drills, about 3 in. apart, and covered lightly. Cultivate well and fertilize. When ready for use, pull every other onion in the row, allowing balance to mature.

All Vegetable Seeds

10c PER PACKET UNLESS OTHERWISE NOTED

In bulk quantities at money-saving prices. Special quotations to market gardeners. Prices subject to change without notice.

Planting Guide FOR VEGETABLE GARDENS


CARROT
Imperator



CABBAGE
Copenhagen Market



SWEET
CORN, *Country Gentleman*



TURNIP, *Purple-Top White Globe*



LETTUCE, *Grand Rapids*



	Seed Required for 50 Ft. of Row	Distance Between Rows (Inches)	Distance Apart in Row (Inches)	Depth of Planting (Inches)	Approximate Yield per 50 Ft. of Row	Seed Required Family of 5
Artichoke	$\frac{1}{8}$ oz.	36 to 48	18 to 24	$\frac{1}{4}$		
Asparagus	$\frac{1}{2}$ oz.	14 to 24	3 to 6	1	25 lbs.	$\frac{1}{2}$ oz.
Beans, Bush	$\frac{1}{2}$ lb.	24 to 30	2 to 3	$1\frac{1}{2}$ to 2	20 qts.	2 lbs.
Beans, Bush Lima	$\frac{1}{2}$ lb.	18 to 24	2 to 3	1	32 qts.	1 lb.
Beans, Pole	$\frac{1}{2}$ lb.	36 to 48	6 to 8	$1\frac{1}{2}$ to 2	30 qts.	1 lb.
Beans, Pole Lima	$\frac{1}{4}$ lb.	36 to 48	6 to 8	$1\frac{1}{2}$ to 2	$1\frac{1}{4}$ bu.	1 lb.
Beets, Table	$\frac{1}{2}$ oz.	14 to 24	3 to 4	$\frac{1}{2}$ to 1	$1\frac{1}{4}$ bu.	2 oz.
Beets, Mangel	$\frac{1}{2}$ oz.	20 to 28	3 to 4	1		
Broccoli, Heading	$\frac{1}{8}$ oz.	24 to 30	24	$\frac{1}{2}$	50 head	1 pkt.
Brussels Sprouts	$\frac{1}{8}$ oz.	24 to 30	12 to 18	$\frac{1}{2}$	15 qts.	1 pkt.
Cabbage	$\frac{1}{4}$ oz.	18 to 36	14 to 24	$\frac{1}{2}$	35 to 50 head	$\frac{1}{4}$ oz.
Chinese Cabbage	$\frac{1}{2}$ oz.	18 to 24	12 to 15	$\frac{1}{2}$	40 to 50 head	1 pkt.
Carrots	$\frac{1}{4}$ oz.	16 to 24	1 to 3	$\frac{1}{2}$	24 qts.	$\frac{1}{2}$ oz.
Cauliflower	$\frac{1}{4}$ oz.	24 to 30	20 to 24	$\frac{1}{2}$	25 to 30 head	1 pkt.
Celery	$\frac{1}{8}$ oz.	24 to 40	4 to 6	$\frac{1}{4}$	75 to 100 stalks	1 pkt.
Collards	$\frac{1}{4}$ oz.	24 to 30	14 to 18	$\frac{1}{2}$	3 bu.	1 pkt.
Corn, Sweet	4 oz.	30 to 42	9 to 12	1	100 ears	1 lb.
Corn, Pop	3 oz.	34 to 40	6 to 8	1		
Cucumber	$\frac{1}{2}$ oz.	48 to 60	12 to 36	$\frac{1}{2}$ to $\frac{3}{4}$	75 to 100 fruit	$\frac{1}{2}$ oz.
Dill	$\frac{1}{4}$ oz.	18 to 36	4 to 8			
Egg Plant	$\frac{1}{8}$ oz.	24 to 36	18 to 24	$\frac{1}{2}$	70 to 80 fruit	1 pkt.
Endive	$\frac{1}{2}$ oz.	18 to 24	8 to 12	$\frac{1}{2}$	50 to 60 plants	1 pkt.
Kale	$\frac{1}{2}$ oz.	24 to 32	14 to 22	$\frac{1}{2}$	50 plants	1 pkt.
Kohlrabi	$\frac{1}{4}$ oz.	14 to 24	4 to 6	$\frac{1}{2}$	1 bu.	1 pkt.
Leek	$\frac{1}{4}$ oz.	12 to 42	2 to 4	$\frac{3}{4}$		1 pkt.
Lettuce	$\frac{1}{4}$ oz.	12 to 18	4 to 12	$\frac{1}{4}$	25 lbs.	2 oz.
Muskmelon	$\frac{1}{2}$ oz.	70 to 80	36 to 60	$\frac{3}{4}$	50 to 60 fruit	1 pkt.
Mustard	$\frac{1}{4}$ oz.	14 to 24	6 to 9	$\frac{1}{2}$	1 to $1\frac{1}{2}$ bu.	$\frac{1}{2}$ oz.
Okra	1 oz.	36 to 48	24	1	450 pods	1 pkt.
Onion	$\frac{1}{4}$ oz.	18 to 24	3 to 4	$\frac{3}{4}$	1 bu.	$\frac{1}{2}$ oz.
Onion (for sets)		12 to 14	not thinned	$\frac{3}{4}$	300 scallions	
Onion sets	1 pt.	12 to 18	2	1	60 bunches	1 qt.
Parsley	$\frac{1}{4}$ oz.	12 to 20	6 to 8	$\frac{1}{2}$	45 bunches	1 pkt.
Parsnips	$\frac{1}{2}$ oz.	18 to 24	3 to 4	$\frac{1}{2}$	1 bu.	$\frac{1}{2}$ oz.
Peas	$\frac{3}{4}$ lb.	24 to 36	1 to 2	1 to 2	30-50 qts. pods	3 lbs.
Peppers	$\frac{1}{8}$ oz.	20 to 30	18 to 20	$\frac{1}{2}$	25 doz. peppers	1 pkt.
Pumpkin	2 oz.	96 to 108	60 to 84	$\frac{3}{4}$	25 to 35 fruit	1 pkt.
Radish	$\frac{1}{2}$ oz.	12 to 18	1 to 2	$\frac{1}{2}$	50 bunches	3 oz.
Rhubarb	$\frac{1}{2}$ oz.	24 to 48	18 to 24			
Rutabaga	$\frac{1}{8}$ oz.	18 to 24	6	$\frac{1}{2}$ to 1		1 pkt.
Salsify	$\frac{1}{2}$ oz.	18 to 24	2 to 4	$\frac{1}{2}$		1 pkt.
Spinach	$\frac{3}{4}$ oz.	14 to 18	3 to 5	$\frac{3}{4}$	1 $\frac{1}{2}$ bu.	1 oz.
Spinach, N.Z.	$\frac{1}{2}$ oz.	30 to 36	12 to 18	1	35 to 50 lbs.	1 oz.
Squash, Summer	$\frac{1}{4}$ oz.	36 to 48	48 (4 seeds)	1	60 to 70 squash	1 pkt.
Squash, Winter	$\frac{1}{4}$ oz.	72 to 90	72 (4 seeds)	1	35 to 40 squash	1 pkt.
Swiss Chard	$\frac{1}{2}$ oz.	12 to 18	4 to 6	1	25 lbs.	1 oz.
Tomato	$\frac{1}{16}$ oz.	40 to 60	36 to 40	$\frac{1}{2}$	2 bu.	1 pkt.
Turnip (roots)	$\frac{1}{4}$ oz.	18 to 24	2 to 4	$\frac{1}{4}$	1 bu.	$\frac{1}{2}$ oz.
Turnip (greens)	$\frac{1}{2}$ oz.	12 to 18	2 to 4	$\frac{1}{4}$	1 $\frac{1}{2}$ bu.	1 oz.
Watermelon	$\frac{1}{2}$ oz.	72 to 96	72 to 96	$\frac{3}{4}$	20 fruit	1 pkt.

VEGETABLE GROWING CHART

ASPARAGUS. Seed should be soaked 24 hours before planting—and covered 2" deep in the garden. Thin young plants to 1" apart. Cultivate often. Set in permanent beds in spring.

BUSH BEANS. Do not plant until danger of frost is past. Press soil firmly around seeds. Thin young plants to about 6" apart. Cultivate only before blooming and never when wet. Pick frequently. Make plantings every two weeks for supplies throughout the season.

POLE BEANS. Plant about same time as Bush Beans. Poles 5' to 8' long should be driven into ground in rows about 4' apart with spacing of 3' in row. Run rows north and south. Plant 5 to 8 beans around each pole, thinning to 4 plants later. Hoe frequently.

Caution: To avoid spreading plant diseases, do not cultivate or pick when plants are wet.

BEETS. Plant about same time as radishes in deep, sandy loam. Have soil fine and loose and press down firmly around seed. When tops are 3" to 6" tall pull them and use for cooked greens. Continue this until roots stand 6" apart. Plant every 10 days or so.

BRUSSELS SPROUTS. Easy to grow wherever conditions are favorable for late cabbage, and requires same culture.

CABBAGE. Set out as soon as frost danger is past. Use plenty of good commercial fertilizer. Cultivate frequently, every 5 or 6 days until cabbages are large. For winter storage, stand plants upright in cold cellar, with roots in sand. Slight freezing will not harm cabbage.

CARROTS. Sow early and thinly—in light, deep soil, preferably fertilized the previous season. Thin plants when about 2" tall. Keep on cultivating frequently throughout season.

CAULIFLOWER. Follow same methods as for cabbage except that heads must be protected from sunlight to assure the desirable white curd. Gather and tie the tops of leaves together loosely to shut off light. Do not cramp the heads.

CHINESE CABBAGE. An easily raised succession crop. Set out in rows which have been occupied by earlier vegetables.

COLLARDS. Sow seed thickly in rich ground and transplant when about 4" high—or if plants are to remain where sown, spread the seed more thinly, and when young plants are well started, space them out to 2' or 3' apart in the row.

CORN. Do not plant until all danger of frost is past. Be sure soil is

well worked and dry—then drop 6 kernels of corn in a shallow hole made with the corner of the hoe. Cover each "hill" with about 1" of fine soil pressed down firmly. When 6" high, thin to three or four plants in each hill. Hoe often, but not deeply.

CUCUMBER. The best soil is warm, sandy loam—preferably on a slight southward slope. Be sure all danger of frost is past, and that soil is fairly moist and loose. Plant 8 or 10 seeds to the hill. When 6" tall, thin to 3 in a hill. Take fruit as soon as it reaches picking stage.

EGG PLANT. Requires continuous warm weather for best results. Seed should be started in a hotbed, as it is slow to germinate. Set plants in open ground when 2" tall and protect from hot sun when young. Be sure to keep young plants developing rapidly. Cultivate freely. To produce large fruits, remove lateral branches so as to reduce number of fruits per plant. Spray to protect from potato bugs.

ENDIVE. Sow at intervals for continuous supply. When well started, transplant or thin to 1' apart. To blanch, tie outer leaves together over the center when plant is nearly grown. Just before killing frosts in fall, dig the plants, taking plenty of soil with roots—pack closely together and store in dark cellar for winter use.

KALE or BORECOLE. Requires moist, well enriched soil. Pick leaves as wanted, or pull whole plant. Leaves are best after a frost.

KOHLRABI. Sow early as possible in light, rich soil. When plants are in the third leaf, thin to 6" apart. Plant at intervals of 10 days for succession of bulbs until hot weather—after which they do not grow. Use while still young and tender, before skin hardens.

LETTUCE. Sow as soon as ground can be worked—or for extra early crops start indoors and transplant when danger of heavy frost is over. Repeat sowing every 2 weeks to insure continuous supply. For leaf lettuce, thin plants to 6" apart. For head lettuce, thin to 8" spacing, in rows 18" apart. Lettuce grows best and heads best in cool, moist weather.

MUSKMELON. Requires long season to develop and is easily injured by frost, or even by cool weather. Before planting, spade in liberal forkful of well rotted manure in each hill. Thin plants to 4 per hill after third leaf develops, and train vines in different directions. Cultivate as long as possible. On moist ground use shingles to hold melons off ground to prevent rotting.

PEPPERS

California Wonder Improved—Number one quality peppers, about 4-in. long and 3½-in. diameter, very smooth, 3 and 4 lobed, glossy green, turning bright crimson when ripe. Remarkably thick walls, tender, sweet. 75 days.

Florida Giant—Considered one of the best shipping varieties in the South. Plants up to 30-in. tall. Vigorous growth and heavy foliage. 76 days.

Long Red Cayenne—The hot favorite. For canning, pickling, drying. Pungent flavored fruit, tapering and twisted. 70 days.

Pimento—Excellent for canning or stuffing. Smooth, heart-shaped, very thick flesh. 73 days.

Red Chili—Low, spreading, 18-in. plant. Extremely pungent fruit. Erect, conical. 82 days.

Ruby King—Abundant, tapering fruit, dark-green, changing to ruby-red. 69 days.

World Beater—One of the best large peppers. Glossy-green, changes to bright red. Mild, very sweet, thick flesh. 75 days.

PUMPKIN

Kentucky Field or Large Cheese—Dull orange, large ribbed fruit. Coarse, deep yellow flesh, good flavor. 120 days.

King of the Mammoth—Largest of all Pumpkins. Valuable for stock feeding and exhibition. Globular, slightly ribbed, flattened, weigh 60 to 90-lb. Flesh solid, yellow-orange, coarse, but good quality. 120 days.

Small Sugar or New England Pie—Desirable for pie or general use. Round, flat end fruit, deep orange. Thick, yellow-orange flesh. 118 days.

Striped Cushaw—For stock feed. Crook-necked, 11-lb., white and green striped, thick, coarse yellow meat. 112 days.

RADISH

Crimson Giant—Large round, 1 to 1½-in. Deep crimson roots. Very crisp and tender for its size. 28 days.

Early Scarlet Globe—Globular, slightly elongated, bright scarlet. 24 days.

Early Scarlet Turnip, White Tipped—Carmine red with white tip. Flesh white, crisp and mild. 26 days.

French Breakfast—Roots thicker toward bottom. 1½-in. long by ½ to ¾-in. Dull scarlet, white tipped, 25 days.

White Icicle—Tapering roots 5 to 5½-in., very white with pure white, mild flesh. 27 days.

Fall and Winter Varieties

Long Black Spanish—7 to 10-in. roots. Black skin, solid white flesh. Late and very hardy. 55 to 60 days.

Rose China Winter—Popular, very distinct winter sort. Good for late salads. Cylindrical blunt, smooth, bright rose-red. Flesh white, very firm and pungent. 52 days.

SALSIFY

Mammoth Sandwich Island—A palatable and nutritious vegetable when cooked. Flavor is remarkably similar to oysters. 120 days.

SPINACH

Bloomsdale Savoy, Long Standing—A few days later than regular Bloomsdale Savoy. Holds from 12 to 14 days longer. 42 days.

Improved or Giant Nobel—Large, vigorous, spreading plant, slow to form seed stalks. Heavy yielder. Large, thick, smooth, deep green leaves. 45 days.

New Zealand—Thrives in dry, hot weather. Can be picked repeatedly all season. Not a true spinach but similar when cooked. 70 days.

Viroflay—Very large, broad thick leaves, slightly crumpled. Old standard variety suitable for spring or fall sowing. Good for canning. 46 days.

SQUASH

Cocozelle (Bush)—Also called Italian Vegetable Marrow. Oblong shaped. Dark green with light green flesh of very delicate flavor. 60 days.

Early Prolific Straightneck—Earlier, smaller, and more productive than Giant Summer Straightneck, and more uniform. Brilliant yellow. Fine for home or market. 50 days.

Early Summer Crookneck—Fruits curved at neck, weigh 3-lb., light yellow, moderately warted. 53 to 55 days.

Early White Bush Scallop—Fruits greenish when young, become white at maturity. 3-in. long, 7½ to 8½-in. wide. 53 days.

Hubbard—Most widely grown of any winter squash. Round, warted, dark green with thick yellow flesh. Weight 12 to 14 lbs. Excellent flavor. 100 days.

Table Queen or Des Moines—Acorn-shaped, green, deeply furrowed; flesh rich yellow, dry, mealy, delicious. Convenient size for serving. 100 days.

Zucchini—Straight, cylindrical fruits, 3 to 4 lb., 4 to 5-in. through, 10 to 14-in. long. Mottled and striped green-creamy-gray. Delicate flavored flesh. Bush type. 60 days.

SWISS CHARD

Lucullus—Most popular Chard. Upright in growth, with yellowish-green curled, crumpled leaves. Thick, broad and light green stems. 50 to 60 days.

TOMATOES

Dwarf Champion—Desirable where space is limited. Plant dwarf tree type, dark green leaves. Fruits medium, smooth, solid, pink, globular. 86 days.

Grothen's Red Globe—Wilt resistant. Deep red with no yellow about stem end. Popular for shipping. 73 days.

Improved Stone—Medium late. Deservedly popular with canners and market gardeners. Ripens evenly and is uniform. Fruits large, flattened, but deep. Smooth, attractive scarlet-red, of fine flavor. 86 days.

Jubilee—Bright orange with few seeds. Good for slicing or juice. Rich in vitamin content. 72 days.

June Pink—Earliest of pink fruited varieties, heavy bearer. Vine open, spreading, rather short. Fruits medium size, flattened, smooth, purplish-pink. 69 days.

VEGETABLE GROWING CHART

OKRA or GUMBO. Do not plant until ground is warm, as this is a tender, hot weather plant. Pick pods before they develop woody fibres.

ONION SETS. Onion sets used instead of seeds will produce earlier crops of green onions or large bulbs. Plant sets right side up and cover with garden rake; then firm the soil well over the sets.

ONION. Plant as soon as soil can be prepared. For best yield, plant on very fertile land—fall plowed, and thoroughly fertilized. When plants are a few inches tall, thin to prevent crowding—using the plants removed as green onions. Those left to become fully ripe can be stored for winter. Cultivate and hand weed crop every 2 weeks during summer.

PARSLEY. Does best in rich, mellow loam. Seed is slow to germinate, and is helped by soaking in warm water over night before planting. Sow early and not too deeply. When curled varieties are about 3" tall, cut off all leaves. The new growth will be brighter and better curled.

PARSNIP. Plant in rich, sandy loam, thoroughly pulverized. Seed requires plenty of moisture for germination and should be sown early. Dig after a killing frost. Freezing improves parsnips, so some can be left in the ground all winter and used in spring. For storage, bury in dry sand.

PEAS. For early crop, plant in light, rich soil. For general crop, a rich loam or clay soil is best. Plant at 2-week intervals for continuous supply. Soak seeds in water the night before planting. Peas need moderate temperature, plenty of moisture. Light frosts do not injure them so they may be planted early in spring. Varieties more than 1½' tall should be staked when 4" to 6" tall.

For increased yield, **INOCULATE!** Garden peas, like all legume crops, almost always benefit by inoculation. Quantity and quality of the crop are increased when the beneficial nitrogen-fixing bacteria are present in sufficient numbers.

PEPPERS. Warm, mellow soil in sheltered location is best. Start under glass. Cultivate regularly, drawing soil up around stems. When plants are 7" to 8" tall, hoe in light dressing of commercial fertilizer. Do not plant hot peppers near sweet; they are apt to cross.

PUMPKIN. Cultivate practically the same as melons or cucumbers.

RADISH. Soil should be light, quick and rich to insure rapid growth. Slow growth makes the

flavor too strong. Sow as early as ground can be worked, as Radishes are very hardy. Make successive sowings up to hot weather. When in the third leaf, thin to give roots plenty of room. Pull promptly when mature. The secret of crisp, delicious radishes is quick growth, prompt picking.

RHUBARB. Sow in cold frame in spring and transplant into rows as soon as plants are large enough. In fall or the following spring transplant to permanent location in rows 3' to 5' apart each way. Stalks should not be taken for use the first year.

SPINACH. Plant very early in spring—or start seed in fall (giving it protection by 3" of straw over the winter) and enjoy an early spring crop. Cut all spinach before hot weather as it doesn't do well in extreme heat. Sow again in August or September for fall crop. Spinach should be gathered before the flower spike appears.

SQUASH. Plant about same time as corn—and in hills. Does best in rich, sandy loam, and is helped by fertilizing the hills. When in the third leaf, thin to 4 plants per hill. Cover every fourth joint with earth to encourage extra root formation. Bush varieties may be planted in hills 3' to 4' apart.

SWISS CHARD. Requires about same treatment as beets. Cultivate frequently. Leaves may be gathered during summer and fall. New ones will grow quickly.

TOMATOES. Do best in sandy, well pulverized loam. Sow seed in hotbed or indoors. When plants are about 2" high, set out, 3" apart, in boxes or pots—later transplant into the garden. Or keep in flats until all danger of frost is past, and then set plants out directly into the garden. Water around the roots of the plants when setting them out, if ground is dry. Cultivate frequently until plants shade the ground. Remove all but the two or three strongest branches. Study methods of staking and training vines and adopt the one best suited to your situation.

TURNIPS. For summer greens or roots, sow as early as radishes and lettuce. Thin out when plants are 1" high. Sow again in summer for fall and winter use. Turnips do best when most of their growth is made in autumn or early spring. Should be grown rapidly for best flavor and texture.

WATERMELON. Requires about same culture as muskmelon, except the vines need more room. Fertilize each hill liberally and cultivate thoroughly.

GARDEN GUIDE POSTS



Growing food at home is just plain good sense. Millions of Americans last year found that vegetables F.O.B. the kitchen door taste better than any they can buy. What's more, scientists tell us that only fresh vegetables contain the all-important vitamins in abundance. That is why most of our home gardeners will continue to grow food at home even though the war emergency may seem to be over.

To help you grow better vegetables and to answer the questions that home gardeners asked during last season, we have completely revised our "GARDEN GUIDE POSTS" for this issue of our catalog.

For Veteran Gardeners

Even old timers will find these notes worth saving for the many items of information that every gardener needs from time to time—when to plant, how deep, how much seed to buy, etc.

For New Gardeners

We've tried to make this a pocket encyclopedia of gardening, with all the basic facts you need condensed in the shortest and simplest form possible. Most of your questions can be answered by referring to the information given below.

SITE AND SOIL

Most of us have no choice in either site or soil and must work with what we have, but if selection is possible, keep these principles in mind:


1. Locate the garden in an open spot that receives at least six hours of direct sunlight each day, away from the roots of trees that might rob the soil of moisture and fertility.

2. A rich sandy loam is the best garden soil, but if you have either heavy clay or light sand, the same materials will improve both—decayed vegetable matter—old rotted manure, peat moss, commercial humus or well-rotted leaves.

3. Don't apply lime unless it is really needed. Usually a soil that will grow good beets is all right. If beets do poorly and grow irregularly, you can add about 50 lbs. of hydrated lime or 100 lbs. of ground limestone to each 1,000 square feet of garden. If you have a pH tester, most flowers and vegetables will do well in a soil between 6.0 and 7.0.

4. Do apply fertilizer. Apply 30 lbs. of good general fertilizer like 4-12-4 or 5-10-5 to every thousand square feet and you won't have to worry about any deficiencies.

5. Don't plant in wet ground. No soil is right for gardening that can't pass the "mud pie" test. To make a mud pie test, pick up a handful of soil and squeeze it slightly. If it balls or packs together and won't crumble when you apply light pressure, it is too wet to work.




8 KEY POINTS IN YOUR PLANNING

1—On level ground, you can run rows either east and west or north and south. If east-and-west layout is used, be sure to plant taller vegetables to the north. On sloping ground, the rows must run across the slope, not up and down. Otherwise all your plant food and much of your topsoil will go down the hill when it rains.

2—Divide the garden into three sections if possible, (1) Root vegetables (2) Leaf vegetables and (3) Fruit vegetables. By switching these three groups every year you avoid many soil-borne diseases and insects and get better use of fertilizers. This isn't absolutely necessary, but it is one of those details that the smart gardener watches.


3—Plan to use your garden full time. To do this, you'll need catch crops, intercrops (also called companion crops) and succession crops. Catch crops are early vegetables, like spinach, that you grow very early before a crop like late cabbage is set out. Intercrops are planted between the rows of larger vegetables, like head lettuce between rows of Broccoli. Succession crops are catch crops in reverse—they follow early crops. For instance, Chinese cabbage will mature fine heads if planted after an early crop of peas has been taken off.




RADISH, Early Scarlet Turnip



BEANS, Golden Wax



BEET, Detroit Dark Red



LETTUCE
Big Boston

TOMATOES

(Continued)

Marglobe — Moderately productive, with heavy foliage. Vigorous and resistant to wilt and rust. Fruits medium large, nearly round, smooth, and solid. Deep scarlet-red. 77 days.

Oxheart — Open, spreading, moderately productive vine. Fruits extremely large, somewhat rough, heart-shaped and pink. Solid, mild-flavored. 90 days.

Pan America, New Wilt-Resistant — Newly developed by the U. S. Dept. of Agriculture. Deep scarlet, heavy and firm. Evenly shaped. Resistant to Fusarium wilt and nailhead rust. Worth a try. 75 days.

Ponderosa — One of the largest varieties. Spreading large vines. Solid, fine flavored, pink, with small seed cells. Somewhat rough and flat. Have a tendency to crack. Late variety. 88 days.

Pritchard — A cross between Marglobe and Cooper's Special, developed by U. S. Department of Agriculture. Large, solid, smooth, round fruit, with thick walls. Rather light scarlet. 73 days.

Rutgers — Second early variety. Large, thick stems, vigorous foliage. Fruits similar to Marglobe but flatter at stem end. Bright red, with thick walls. Red, firm flesh, low acidity. 72 days.

Stokesdale — An extremely important new development. Smooth, 7-oz. fruit, brilliant color, exceedingly solid flesh and small seed cavities. Matures in the second-early group, a full week earlier than the Marglobe group. Usually ahead of Bonny Best, too—and 2 ounces heavier.

Small Fruited

Red and yellow varieties, shaped like miniature pears, plums, and cherries; very attractive in appearance and popular for preserves.

TURNIPS

White Fleshed

Amber Globe — Chiefly grown for stock, but used for table when young. Tops large, cut leaved. Roots round, large, smooth, globe-shaped, light yellow with green top. Flesh yellowish white, of good quality. 76 days.

Long White Cowhorn — Long white variety used principally for stock food. Tender and sweet for table use when young. Tapered; 12-15 in. long. 65 days.

Purple Top — All purpose variety. Large, sweet tops. Smooth, globe-shaped roots, purple-red above, white below. Sweet, tender, crisp, fine grained white flesh. 50 to 60 days.

White Egg — Cut leaved, medium sized, upright tops. Egg shaped roots, all white. Smooth grained. 55 to 60 days.

Foliage Turnip

Seven Top — For greens or forage. Young shoots extremely tender. Roots tough, undesirable for food. 50 days.

Shogoin or Japanese Foliage — Recently introduced and rapidly gaining popularity. 18 to 20-in. bright green tops, strap leaved. Roots are semi-round, and white. Very mild and tender. 70 days.

Rutabaga or Swede Turnip

American Purpletop — Hardy and productive. Tender firm, light yellow, sweet flesh. 88 days.

(Cont'd from preceding page)

4—Avoid growing vegetables your family doesn't like and won't eat, but be sure to include plenty of those they do like. If they're fond of beans, for instance, don't be satisfied with a single large planting, but make several smaller plantings at two weeks intervals.

From thousands of reports received from home gardeners on last year's results, Tomatoes, Swiss Chard, Green Beans and Carrots produced the most food with the least effort. Concentrate on these if your space is very limited.

5—in smaller gardens, don't plant space-wasting crops like Potatoes, Pumpkins, Sweet Corn and Vine Squash. On the farm these can be grown with the field crops, saving the more highly cultivated garden space for the more productive vegetables that need closer attention.

Many gardeners last year reported that Zucchini and other bush squashes gave better results than Eggplant and satisfied many of the requirements for vining Squashes.

6—Include perennial vegetables when possible. Rhubarb, Asparagus, Horse Radish and Perennial Onions can be grown on one side of the garden where they won't interfere with plowing or cultivating. Small fruits should be included whenever space will permit, since the quality of home-grown berries is far better than any you can buy.

7—Plan to grow enough vegetables for storing and canning. Remember that correct storage begins with planting (see under Storage) so provide for this when planning your garden and ordering seed.

8—Study the chart on page 15. This will tell you how far apart rows should be, how long each crop takes to mature, and how much space you will need to provide the vegetables your family likes. After you have taken into consideration the crops you want to grow, make a rough sketch showing the location of each crop, with catch crops, intercrops, succession crops indicated.



MAKE A PERMANENT PLAN

After the rough sketch has been made, this information should be transferred to a larger sheet with the garden drawn to scale.

A scale of one-quarter of an inch representing one foot works very well, because by using 1/16 of an inch as three inches, almost any standard spacing of the rows can be shown in exact scale. Save this plan from year to year so you can remember where each crop was. This will enable you to rotate root crops, leaf crops and fruit crops.

With the plan complete, you are now ready to order your fertilizer, seed, tools and insecticides. It is to your advantage to do this as early as you can. Late in the season many desirable varieties may be sold out, deliveries may be slow and clerks are so busy that they can't help you as well.

STARTING SEED INDOORS

Many plants can be started indoors or in coldframes or hotbeds outdoors for earlier crops.

Any gardener can do this, but it cannot be done carelessly if good plants are to be grown. If care cannot be given, it would be better to sow crops like Tomatoes and Cabbage direct out of doors because direct-seeded plants will produce bigger crops than will transplants that have been damaged by high heat and too little water.

Tomatoes, Cabbage, Cauliflower, Broccoli, Peppers and Kohlrabi can be started indoors. Use a box three inches deep and of any suitable size. Space the bottom boards about an eighth of an inch apart to provide drainage. Sift soil and use the coarser part in the bottom, with an inch of fine soil on top. Press around edges of box with a brick or board to firm the soil. Add more sifted soil and draw the straight edge of a lath or ruler across the top to level.

Press furrows about one-quarter of an inch deep into the soil, the furrows spaced about 2" apart. By treating the seed with Rootone, a plant hormone that speeds up germination and increases the strength of seedling roots, better seedlings which will transplant with less loss can be grown. Plant seeds 3/4" apart to allow room to lift each seedling without disturbing the rest. After seeding, fill furrow with clean sand or sifted soil and press down soil over entire box until it is 1/2" below upper edge.

WATERING AND TEMPERATURE

Use either a small fly sprayer or a rubber spray bulb that throws a fine mist to water the box after planting. If not available, cover surface with burlap and water through this.

Correct control of moisture is the most important single factor in success with indoor seed sowing. Soil should never be saturated, yet must not dry out at any time until seedlings show second set of leaves, when they can be transplanted outdoors if weather permits.

If seeds of several varieties are grown together, keep box at 70 degrees until seeds germinate, but if only tomatoes and peppers are grown, a temperature of between 75 and 85 degrees will produce better seedlings.

Don't start seedlings too soon or they may become long and spindly. Six weeks before time to set out is soon enough to sow seed indoors, and eight weeks if a hotbed or coldframe is used.

Answers to QUESTIONS THEY ASK US

How long should seed be soaked before planting?

No time at all, except in hot dry weather, when it sometimes helps to speed germination by soaking some seed, but never for more than 12 hours.

In the spring, when moist soil is overly wet anyway, nothing is gained by soaking seed, and often damage is done. Seeds, like people, expire from oxygen starvation when held under water too long.

Doesn't squash take up more room than it's worth?

Even the smallest garden should in-

clude some of the bush types of squash, such as Zucchini, and Prolific Straight-neck. They don't need much space, are easy to grow, and give lots of vegetables for little trouble. These small varieties are delicious eating, and can be a new taste treat when baked with cheese, or fried like eggplant.

However, unless your garden is large, stay away from the vine type squash, as it is usually not desirable for growing in small areas.

My carrots always do poorly. Why?

In almost every case, this is because the seed was sown too thickly, and not

thinned out. Never leave more than one seedling for every 2 or 3 inches of row, and never sow the long, slender varieties of carrots in heavy clay soil.

In growing sweet corn, what are the advantages of hybrid varieties over the old fashioned types?

Hybrid sweet corn has larger ears, deeper kernels, and is more resistant to certain diseases than the old-time varieties. Too, all the ears mature at the same time, so there is never any doubt as to which ears should be picked first. The stalks are stronger, and hybrid varieties produce more corn per square feet of planting space.

Why don't I get crisp radishes?

The secret of getting crisp, delicious radishes is quick growth, so use plenty of plant food in the soil, where they are planted. Radishes are the easiest of all vegetables to grow, as the seeds germinate in a few days, and are ready for your table in 3 to 5 weeks. Sow a succession of radish seed every 10 days all through the planting season—grow them quickly and you will have plenty of crisp, tender radishes.

I never have any luck getting summer-sown seed to grow. What's wrong?

The biggest problem here is because most summer-sown seed are those of cool-weather crops that we want to mature when the cool days come, yet the seed must be started during high summer temperatures.

For success with summer-sown seed, first try chilling the seeds in your refrigerator for several days.

Then, plant the seeds in soil containing lots of decayed vegetable matter so it will hold all the moisture you can pour on. As the moisture evaporates from the surface, it will cool the soil.

If your garden plot is exposed to dry, hot winds, planting summer-sown seed to the lee side of tall plants, such as tomatoes, corn, etc., will keep the dry air from searing the tender seedlings.

After sowing, wet the rows with cool water, and cover with boards until the seeds have sprouted. Remove boards late in the day, or during cloudy weather to prevent the tender seedlings from sunscald.

I have clay soil in my garden. What can I grow?

When modified with well-rotted manures or humus, clays are rated with our best soils. However, if the clay is stiff and hard, successful growth of root crops, like beets, carrots, turnips and parsnips, is especially difficult, and others are handicapped, too.

What grows in sandy soil?

You can grow practically everything if the soil is good sandy loam—and this type of soil is perhaps the finest for vegetables. However, sandy soils need plenty of plant food.

Be cautious, though, if the soil is mostly sand, as little will grow here.

KNOW YOUR GARDEN ENEMIES!

To find the right treatment for insect enemies, identify them in the "Rogues' Gallery"


below. Then refer to our free Spray Guide Chart for proper treatment in each case.



Cutworms



White Grub



Chicken Louse



Mexican Bean Beetle



Rose Mildew



Coddling Moth



Potato Blight



Rose Aphis



Tomato Worm



Potato Bug



White Flies



Sow Bug



Rose Leaf Spot



Rose Midge



Asparagus Beetle



Japanese Beetle



Lace Wing



Potato Flea Beetle



Thrip



Rose Chafer



Mealy Bugs



Orthezia



Slugs



Caterpillar



Leaf Hopper



San Jose Scale



Red Spider



Oyster Shell Scale



Aphids



Tarnish Bug

Which vegetables are actually the best for high vitamin content?

All fresh vegetables are high in the vitamins which we all need for proper nourishment. If we could eat enough vegetables fresh from the garden, we wouldn't need to buy our vitamins in pills at the drug store.

But some vegetables are especially high in certain vitamins—

For Vitamin A—eat plenty beet greens, broccoli, collards, endive (escarole), mustard greens, and spinach.

For Vitamin B—brussels sprouts, kale, green lima beans, and peas.

For Vitamin C—beet greens, broccoli, brussels sprouts, cauliflower, kale, mustard greens, peppers and spinach.

For Vitamin G—beet greens, collards, kale, mustard greens and spinach.

What are some good combinations for succession planting? For "intercropping"?

The early crops that can be harvested in time to be followed by others are: Early cabbage, cauliflower, lettuce, peas, beets, radishes, carrots, kohlrabi, turnips, spinach, and green onions from sets.

These crops may be followed by:

Peas (in cool sections only), late celery, late plantings of lettuce, beets, carrots, turnips, cabbage, cauliflower, and early varieties of sweet corn.

For late autumn and winter vegetables, plant chinese cabbage, kale, mustard, after your early crops are harvested. For intercropping, plant rows of quick growing items such as lettuce, radishes, or spinach between the wide spaced rows of slower growing vegetables, such as celery, peas and tomatoes.

Set early cabbage between the rows where tomatoes are to be planted later, and bush beans between rows where eggplant and peppers will follow.

How can I get rid of crab grass?

Crab grass doesn't start until warm weather, so your very first step in fighting this lawn enemy No. 1, is to plant plenty of good seed early, and fertilize early. When your lawn has a vigorous start, it doesn't give the weed an opening.

Next, since crab grass is an annual which propagates itself by re-seeding, treat your lawn with lead arsenate 20-25 pounds per 1000 square feet before last year's crab grass seed has a chance to sprout. Apply this as a dust or spray. Damage to the blue-grass will be avoided if this treatment is followed by application of a fertilizer rich in nitrogen.

Where crab grass has actually appeared, a direct chemical attack is necessary. New weed killers have been perfected which kill the pest without injury to your lawn. (Ask us about them.)

Along with these methods of attack, you can regulate your ordinary routine of mowing, watering, and fertilizing to help the fight on crab grass. High cutting helps, because a thick mat of grass has a better chance to smother the crab grass seedlings. Watering infrequently but heavily helps, too, because the young seedlings can't reach the deep moisture.

If you want to PLANT BY THE MOON

There is quite a lot of discussion about the effect of the moon on various crops. While we make no attempt to take one side or the other on this question, we are giving all the necessary information in condensed form.

Naturally, we can't vouch for the effectiveness of this method of planting. We are simply submitting the up-to-date charts, etc., for gardeners who observe the phases of the moon when planting.

Signs of the Zodiac and Their Influence

♈ ARIES: Tends to dryness, barrenness. Fairly good for cultivating, spraying and destroying weeds. Plant onions.

♉ TAURUS: Fairly productive. A good sign for planting root crops, particularly potatoes and bulbous plants.

♊ GEMINI: Dry and barren. Not good for any kind of planting or transplanting. Kill noxious growths.

♋ CANCER: Moist, very fruitful, most productive of the Zodiac. Planting and transplanting should yield good crops. Irrigate, bud, graft, transplant.

♌ LEO: Most dry and barren sign. Use only to kill weeds, destroy roots, etc.

♍ VIRGO: Moist but barren. Not recommended for planting or transplanting; good for cultivating and destroying.

♎ LIBRA: Fairly fruitful. Seed for hay, corn, fodder, etc. Produces vigorous pulp growth, reasonable amount of grain. Good for flowers.

♏ SCORPIO: Next to Cancer in productiveness. Irrigate but do not dig potatoes.

♐ SAGITTARIUS: Generally considered barren but used by many for onions, seeding for hay, destroying weeds, etc.

♑ CAPRICORN: Earthly, fairly productive. Similar to Taurus but drier. Bulbous flowers, potatoes, tubers, root crops.

♒ AQUARIUS: Somewhat barren. Good only for cultivation and extermination of pests.

♓ PISCES: Similar to Cancer so use for same purposes. Very productive. Do not dig potatoes.

When to Plant

The table at the right shows when the moon enters each sign of the Zodiac. It also shows the moon's changes.

Under each month, we show first the time at which the moon enters each SIGN (day of the month, and hour of the day, Central Standard Time). Then we show the moon CHANGES (day and hour) for each month.

Ⓐ 1st QUARTER, MOON INCREASING—During 1st Quarter of Moon plant the following: Asparagus, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Corn, Cucumbers, Endive, Kohlrabi, Leek, Lettuce, Spinach, Barley, Oats, Wheat, Rye.

Ⓑ 2nd QUARTER, MOON IN-

1945—Moon Signs and Changes—1945

JANUARY	APRIL	JULY	OCTOBER		
2 ♑ 10:52 a.	1 ♈ 9:06 p.	1 ♉ 6:29 p.	2 ♑ 11:36 a.		
4 ≈ 10:45 p.	4 ♀ 7:48 a.	3 ♈ 9:04 p.	4 ≈ 10:17 p.		
7 ♓ 11:11 a.	6 ≈ 3:26 p.	5 ♉ 11:20 p.	7 ♓ 10:26 a.		
9 ♂ 9:53 p.	8 ✶ 7:09 p.	8 ☽ 2:13 a.	9 ♂ 11:16 p.		
12 ♀ 5:24 a.	10 ♈ 7:37 p.	10 ☽ 6:48 a.	12 ♀ 11:30 a.		
14 ≈ 9:54 a.	12 ♀ 6:39 p.	12 ♑ 2:01 p.	14 ≈ 9:03 p.		
16 ✶ 12:26 p.	14 ☿ 6:32 p.	15 ≈ 0:14 a.	17 ✶ 2:28 a.		
18 ♈ 2:22 p.	16 ☽ 9:17 p.	17 ♓ 0:29 p.	19 ♈ 4:07 a.		
20 ♀ 4:49 p.	19 ☽ 3:57 a.	20 ♂ 0:33 a.	21 ♀ 3:30 a.		
22 ✶ 8:35 p.	21 ♑ 2:05 p.	22 ♀ 10:25 a.	23 ✶ 2:54 a.		
25 ☽ 2:08 p.	24 ≈ 2:16 a.	24 ≈ 5:15 p.	25 ☽ 4:16 a.		
27 ♀ 9:36 a.	26 ☿ 5:52 p.	26 ✶ 9:26 p.	27 ♀ 9:00 a.		
29 ♑ 7:09 p.	29 ✶ 2:55 a.	29 ♈ 0:07 a.	29 ♑ 5:12 p.		
CHANGES		31 ♀ 2:29 a.	CHANGES		
● 6 6:47 a.	● 5 1:18 p.	● 2 0:13 p.	● 11 2:22 p.		
● 13 11:06 p.	● 12 6:29 a.	● 9 7:35 a.	● 14 3:38 a.		
● 20 5:48 p.	● 19 1:46 a.	● 17 1:01 a.	● 20 11:32 p.		
● 28 0:41 a.	● 27 4:33 a.	● 24 8:25 p.	● 27 4:30 p.		
FEBRUARY		● 31 4:30 p.	NOVEMBER		
1 ≈ 6:48 a.	1 ♀ 1:39 p.	1 ≈ 4:09 a.	3 ♈ 4:29 p.		
3 ♓ 7:22 p.	3 ≈ 10:03 p.	6 ≈ 5:18 a.	6 ♂ 5:36 p.		
6 ♂ 6:54 a.	6 ✶ 3:16 a.	4 ☽ 9:25 a.	8 ✶ 2:46 p.		
8 ♀ 3:29 p.	8 ♈ 5:21 a.	6 ☽ 4:29 a.	11 ≈ 3:54 a.		
10 ≈ 8:11 p.	10 ♈ 5:28 a.	8 ♑ 10:26 p.	13 ✶ 11:01 a.		
12 ✶ 9:51 p.	12 ✶ 5:16 a.	11 ≈ 8:23 a.	15 ♈ 2:21 p.		
14 ♈ 10:13 p.	14 ☽ 6:57 a.	13 ♓ 8:26 p.	17 ♈ 2:46 p.		
16 ♀ 11:07 p.	16 ☽ 0:01 p.	16 ♂ 8:54 a.	19 ✶ 2:04 p.		
19 ✶ 2:05 a.	18 ♑ 8:57 p.	18 ✶ 7:30 p.	21 ☽ 2:17 p.		
21 ☽ 7:46 a.	21 ≈ 8:44 a.	21 ≈ 2:28 a.	23 ♈ 5:13 p.		
23 ☽ 4:00 p.	23 ♈ 9:21 p.	23 ✶ 6:02 a.	26 ♈ 0:03 a.		
26 ♑ 2:15 a.	26 ✶ 9:09 a.	25 ♈ 7:30 a.	28 ≈ 10:20 a.		
28 ≈ 1:57 p.	28 ✶ 7:23 p.	27 ♈ 8:36 a.	30 ♈ 10:42 p.		
CHANGES		31 ☽ 3:02 p.	CHANGES		
● 5 3:55 a.	● 5 0:02 a.	● 7 6:32 p.	● 4 5:11 p.		
● 12 11:33 a.	● 11 2:21 p.	● 15 6:26 p.	● 12 5:34 p.		
● 19 2:38 a.	● 18 4:12 p.	● 23 6:03 a.	● 19 9:13 a.		
● 26 6:07 p.	● 26 7:49 p.	● 29 9:44 p.	● 26 7:28 a.		
MARCH		● 29 9:44 p.	DECEMBER		
3 ♓ 2:32 a.	2 ✶ 9:21 a.	3 ♈ 9:20 p.	3 ♂ 11:29 a.		
5 ♂ 2:43 p.	4 ♈ 1:01 p.	5 ♑ 5:39 a.	5 ✶ 11:22 p.		
8 ✶ 0:33 a.	6 ♈ 2:22 p.	7 ≈ 3:49 p.	8 ≈ 9:31 a.		
10 ≈ 6:34 a.	8 ✶ 3:15 p.	10 ♈ 3:48 a.	10 ✶ 5:19 p.		
12 ✶ 8:46 a.	10 ☽ 5:03 p.	12 ♂ 4:37 p.	12 ♈ 10:12 p.		
14 ♈ 8:32 a.	12 ☽ 9:22 p.	15 ✶ 4:07 a.	15 ♈ 0:27 a.		
16 ♀ 7:58 a.	15 ♑ 5:12 a.	17 ≈ 0:15 p.	17 ✶ 1:03 a.		
18 ✶ 9:10 a.	17 ≈ 4:06 p.	19 ✶ 4:17 p.	19 ☽ 1:25 a.		
20 ☽ 1:35 p.	20 ♈ 4:35 a.	21 ♈ 5:10 p.	21 ☽ 3:36 a.		
22 ☽ 9:34 p.	22 ♂ 4:27 p.	23 ♈ 4:53 p.	23 ♈ 8:49 a.		
25 ♑ 8:13 a.	25 ✶ 2:11 a.	25 ✶ 5:32 p.	25 ≈ 5:44 p.		
27 ≈ 8:15 p.	27 ≈ 9:34 a.	27 ☽ 8:40 p.	28 ♈ 5:44 a.		
30 ♈ 8:51 a.	29 ✶ 2:50 p.	30 ☽ 2:51 a.	31 ♂ 6:32 p.		
CHANGES		30 ☽ 2:51 a.	CHANGES		
● 6 10:30 p.	● 3 7:15 a.	● 6 7:43 a.	● 4 0:06 p.		
● 13 9:51 p.	● 9 10:26 p.	● 14 11:38 a.	● 12 5:05 a.		
● 20 1:11 p.	● 17 8:05 a.	● 21 2:46 p.	● 18 8:17 p.		
● 28 11:44 a.	● 25 9:08 a.	● 28 5:24 a.	● 26 2:01 a.		
JUNE		SEPTEMBER			
2 ✶ 9:21 a.	7 ≈ 3:49 p.	3 ♈ 9:20 p.	5 ✶ 11:22 a.		
4 ♈ 1:01 p.	10 ♈ 3:48 a.	5 ♑ 5:39 a.	8 ≈ 9:31 a.		
6 ♈ 2:22 p.	12 ♂ 4:37 p.	7 ≈ 3:49 p.	10 ✶ 5:19 p.		
8 ✶ 3:15 p.	15 ✶ 4:07 a.	10 ♈ 4:17 p.	12 ♈ 10:12 p.		
10 ☽ 5:03 p.	17 ≈ 0:15 p.	12 ♈ 5:10 p.	15 ♈ 0:27 a.		
12 ☽ 9:22 p.	19 ✶ 4:17 p.	21 ♈ 4:53 p.	17 ✶ 1:03 a.		
14 ☽ 10:11 a.	21 ♈ 5:10 p.	23 ♈ 5:32 p.	19 ☽ 1:25 a.		
16 ☽ 2:11 a.	25 ✶ 8:40 p.	25 ✶ 8:40 p.	21 ☽ 3:36 a.		
18 ✶ 9:34 a.	30 ☽ 2:51 a.	30 ☽ 2:51 a.	23 ♈ 8:49 a.		
20 ☽ 2:50 p.	31 ♂ 6:32 p.	31 ♂ 6:32 p.	25 ≈ 5:44 p.		
CREASING		CHANGING			
During 2nd Quarter of Moon plant the following: Beans, Egg Plant, Melons, Peas, Pepper, Pumpkin, Squash, Tomatoes, flowers, and cereals.					
When possible, plant seed while the Moon is in the fruitful signs Cancer, Scorpio or Pisces. The next best signs are Taurus, Libra and Capricorn.					
During 3rd Quarter, Moon DECREASING—In 4th or Last Quarter of the Moon turn sod, pull weeds and destroy noxious growths, especially when Moon is in the barren signs, Gemini, Leo, Virgo or Aquarius. If you must plant in 4th quarter, be sure to use the dates given in "Best Planting Dates."					

CREASING—During 2nd Quarter of Moon plant the following: Beans, Egg Plant, Melons, Peas, Pepper, Pumpkin, Squash, Tomatoes, flowers, and cereals.

When possible, plant seed while the Moon is in the fruitful signs Cancer, Scorpio or Pisces. The next best signs are Taurus, Libra and Capricorn.

During 3rd Quarter, Moon DECREASING—In 4th or Last Quarter of the Moon turn sod, pull weeds and destroy noxious growths, especially when Moon is in the barren signs, Gemini, Leo, Virgo or Aquarius. If you must plant in 4th quarter, be sure to use the dates given in "Best Planting Dates."

Growing Notes FOR BULB PLANTERS

GLADIOLUS

Make small plantings of Gladiolus corms every 10 days from the time Narcissus are in bloom until last flowers on the Peonies and Irises have faded.

Soil must be well-drained. Water if soil is not moist during summer droughts. In heavy soils, plant at bottom of 5 in. deep trench, in light soils, at bottom of 7 in. deep trench. Set in bulbs and cover with 1 in. of soil, gradually filling in trench as shoot grows until soil is level.

If soil is fertilized as described under Vegetable Garden section, no further feeding will be needed. Stake stems if straight spikes are needed and wind is strong.

Storing Glad Bulbs

Bulbs can be dug for storage seven weeks after last flower fades, but the longer digging can be delayed the better the bulb will keep, unless fall rains may start up growth after a summer dry spell. Bulbs must be dug before ground freezes. Store as close to 33 degrees as possible until February 1st. Then bring into warm room (above 55 degrees) for two weeks and return to 33 degree storage until planting time. This cold-warm-cold treatment will kill all hibernating thrips.

Thrips do come in from outside sources, however. Spray plants from the time shoots show until flower spike appears with following spray: 1½ teaspoons tartar emetic, ½ cup brown sugar or honey and 1 gal. water. Apply every 8 to 10 days.

DAHLIAS

Plant single tubers, each with eye or shoot, as clumps make only mass of foliage and poor flowers. Plant at corn planting time, setting tubers at bottom of 6 in. hole and covering with 5 in. of soil, leaving slight depression to catch moisture.

Dahlias need well-drained moist soil and watering during dry spells. Use same fertilizers recommended under Vegetable Garden section. When first flowers are in full bloom, apply a cupful of sulphate of potash (or a double handful of wood ashes) to each plant, plus a cupful of superphosphate, working this mixture into soil.

Dig clumps after a killing frost and store below 40 degrees, but above freezing. In spring wait until sprouts show and divide into single tubers, being careful to include one eye with each tuber.

BULBS AND TUBERS for Spring Planting

DAHLIAS

Avalon—Fine form, bright golden-yellow. Blooms freely.

Jersey Beauty—Supreme among pink Dahlias.

Jane Cowl—Deep, massive blooms, buff and old gold petals curled and twisted.

Mrs. I. De Ver Warner—Large blooms, old rose suffused lilac.

Jersey Beacon—Chinese scarlet, lighter reverse. Long stems.

Thomas A. Edison—Gorgeous blooms of rich royal purple.

Kentucky—Salmon-pink with yellow and gold blend.

Monmouth Champion—Huge brilliant orange flowers, perfect stems.

White King—Magnificent large blooms, pure snow-white.

Pompon Dahlias—Charming "button" dahlias that require little care but reward you with gorgeous, cut-flowers summer and fall. In mauve, pinks, red, white, bronze, yellow.

CANNAS

King Humbert—Gigantic flowers, orange-scarlet, with bright red markings. Bronze leaved. 5 feet.

Yellow King Humbert—Golden-yellow, dotted red. Green foliage. 4 feet.

President—The best red Canna. Immense flowers of rich scarlet. Green foliage. 4 feet.

Hungaria—The best true pink Canna. Large flower trusses; green foliage. 3½ feet.

Wintzer's Colossal—Largest flowering. Vivid, blazing scarlet. Green foliage. 5 feet.

CALADIUM

Fancy-leaved Caladium—Few summer foliage plants can equal it for rich, beautiful color and dramatic effect.

Elephant's Ear—One of the largest ornamental plants for beds and borders. Huge green leaves give unusual effect.

TUBEROSES

One of the most fragrant of all flowers. Tall spikes of double waxy blooms borne on tall stems. Spear-shaped, grass-like foliage. Annual replacement frequently advisable.

GLADIOLUS

Bagdad—Smoky old rose, lighter throat. Very large. 6 to 8 florets on tall, straight spike. Late.

Beacon—Brilliant, clear, dark rose, with a pronounced blotch. Florets slightly waved and ruffled. One of the best.

Betty Nuthall—Coral-pink, yellow throat markings. Florets of fine texture, large, strong spikes.

Bit o' Heaven—Medium-sized florets of orange with a yellow throat. Tall and straight with 9 to 12 flowers.

Commander Koehl—Fine red. Tall spike. Florets very large, dark scarlet without markings.

Charles Dickens—Bright, radiant purple. Many blooms open at a time, perfect placement.

Dr. F. E. Bennett—Fiery orange-scarlet. Very large flowering.

Debonair—LaFrance pink, shading into shrimp pink, creamy throat blotch. 6-8 medium size blooms.

Golden Dream—Clear, golden-yellow flowers, medium size, tall stems, many open at a time. Late.

Minuet—Beautiful, clear light lavender. Considered the best lavender. Very large flower of heavy substance.

Mother Machree—Smoky lavender overlaid at edge with salmon-pink. 8-12 medium size blooms. Late.

Maid of Orleans—Milky white, with light cream throat. Tall spike, 6-8 perfectly placed florets.

Picardy—Rich apricot-pink. Florets extremely large, slightly ruffled and of wax-like appearance.

Pelegrina—Dark blue, 6 or more well placed blooms open with most of remaining buds showing color. Early. Best dark blue.

Red Phipps—Unusual shade of light red. Medium tall, long flower heads. Late mid-season.

Rosa van Lima—Pure self-colored light pink with many flowers open at one time. Strong, straight stem.

Shirley Temple—Newly developed prize winner. Enormous creamy white flower with deeper cream blotch on lower petals. Tall and lovely.

Vagabond Prince—Iridescent mahogany brown, lighter in throat. Small scarlet blotch on lip. 8-10 medium size blooms open at a time.

Wasaga—Clear glowing buff with 4-5 flowers open at a time. Ruffled. Late mid-season.



Bulbs and tubers are a good INVESTMENT

Few garden flowers pay such high dividends as bulbs and tubers!

Lasting for years...they return your investment over and over...by giving so much beauty for such little care, or cost, or effort.

CANNA
Hungaria



DAHLIA, Jersey Beacon




FANCY LEAVED CALADIUM
— The best summer foliage plant for rich, beautiful color and dramatic effect.


Flower Seed DEPARTMENT



AQUILEGIA



ASTER
American Branching



CENTAUREA
Jubilee Gem

KEY TO SYMBOLS

The symbols after the name of each flower give you the following information:

- (a)—Annual
 - (b)—Biennial
 - (hp)—Hardy Perennial
 - (hhp)—Half-hardy Perennial
 - Suited to full shade
 - ◆—Suited to semi-shade
- For key to cultures see page 31.

AGERATUM (a)

Culture II. Cover $\frac{1}{8}$ -in. Ht. various. Blooms early summer to frost.

Blue Cap—New, dwarf. Rich, deep blue. 6-in. plants. Pkt. 15c.

Blue Perfection—Deep amethyst-blue. 9-in. plants.

Midget Blue—1940 All-America. Compact 2 to 3-in. plants. Azure blue. Pkt. 25c.

ALYSSUM (a) ♦

ANNUALS. **Culture II.** Cover $\frac{1}{4}$ -in. Ht. various. Blooms early summer to frost.

Carpet of Snow—White flowers. Plants about $1\frac{1}{2}$ ft. across and only 3-in. high.

Little Gem—Clouds of honey-scented white flowers.

Lilac Queen—Lavender-lilac flowers, 6-in. plants.

Violet Queen—Dwarf, compact, free flowering. Small, sweetly scented flowers; rich, deep shade of violet. Pkt. 15c.

ALYSSUM (hp) ♦

PERENNIALS. **Culture VII.** Cover $\frac{1}{8}$ -in. Ht. 3 to 9 in. Blooms early spring.

Saxatile Compactum (Gold Dust)—Brilliant golden-yellow flowers and grayish-green foliage on compact, spreading plants. Favorite for rock gardens. 12 in.

ANCHUSA, Blue Bird (a) ♦—**Culture II.** Cover seeds $\frac{1}{8}$ -in. Sprays of indigo blue flowers; 20 to 24-in. compact plants.

ANTIRRHINUM (a) ♦

(Snapdragon)

Culture I. Cover $\frac{1}{8}$ -in. Ht. various. Sow early. They germinate slowly.

Maximum (Tallest Varieties)—Rust-resistant. Best for wealth of flower spikes. Mixed. Pkt. 15c.

Large Flowered (Tall Giant)—A fine strain for bedding or cutting as the

spikes are heavy and long. Mixed. Pkt. 15c.

Super Majestic (Semi-tall)—Large flowers on semi-tall plant. Rust-proof. A fine mixture of colors in this type. Mixed. Pkt. 15c.

Half-Dwarf (Large Flowered)—Very good for bedding. Heavy spikes and good length. Large blooms. Mixed. Pkt. 15c.

AQUILEGIA (Columbine) (hp) ♦—**Culture VII.** Cover $\frac{1}{8}$ -in. Ht. $1\frac{1}{2}$ to 2-ft. Prefers light shade, fairly rich soil.

Mrs. Scott Elliott's Long-Spurred—Extremely large flowers, wide range pastel colors. Pkt. 20c.

ARCTOTIS (Blue-eyed African Daisy) (a)—**Culture I.** Cover $\frac{1}{8}$ -in. Ht. $2\frac{1}{2}$ -ft. Midsummer to frost. Prefers sun.

Grandis—Pure white blooms, bluish center with gold band.

ARMERIA (Thrift) (hp)—Rock-garden or edging plant of bright rose or rose and white. Clustered flowers. Pkt. 15c.

ASTER (a)

Culture I. Cover $\frac{1}{8}$ -in. Ht. various. Need rich soil, good sun.

American Branching—Most popular. $3\frac{1}{2}$ to 4-in. double blooms. Branching. Plants 2 to 3-ft. Long stems. All colors.

California Giant Improved—5-in. double flowers, with long curled petals; stems $1\frac{1}{2}$ to 2-ft. long. Branching 2 to 3-ft. plants. All colors. Pkt. 15c.

Giant Crego (Wilt-resistant)—Flowers 4 to 5-in. double, with ribbon-like petals curled and twisted. Vigorous, branching 2-ft. plants. Long stems. All colors. Pkt. 15c.

Giant Crego Navy Blue—1943 All-America. Large 4-in. long petaled double flowers, deep purple blue, almost blackish blue. True, uniform, 12 to 20-in. branching plants. Pkt. 25c.

Queen of the Market (Wilt-resistant)—First to bloom. Nearly round, double, 3-in. flowers, long stems. Plants 18-in., spreading. All colors.

BABY'S BREATH—See Gypsophila.

BACHELOR'S BUTTON (a) (Centaurea or Cornflower)

Culture IV. Cover $\frac{1}{4}$ -in.

Blue Boy—Very fine, fully double. Blue. Pkt. 15c.

Jubilee Gem—All-America 1937. Compact 1-ft. plants, completely covered with large, double, deep blue flowers. Pkt. 15c.

Pinkie—Double, deep pink. Pkt. 15c.

Red Boy—Deep glowing red. Pkt. 15c.

White—Glistening snow-white.

Mixed—

BALLOON VINE (Love-in-a-Puff) (a)—**Culture III.** Cover $\frac{1}{2}$ -in. Ht. 8 to 10-ft. Prefers warm location. Rapid-growing climber with white flowers, succeeded by small, balloon-like seed-pods.

BALSAM (Lady-Slipper) (a) ♦—**Culture II.** Cover $\frac{1}{4}$ -in. Ht. 2-ft. Double, camellia-like flowers. Wants partial shade, plenty of moisture. Sow late.

BLUE LACE FLOWER (Didiscus) (a)—**Culture I.** Cover seeds $\frac{1}{8}$ -in. Ht. 2-ft. Blooms midsummer. Clusters of lavender-blue $2\frac{1}{2}$ -in. flowers, like tiny parasols.

BURNING BUSH or FIRE BUSH—See Kochia.

CALENDULA (a) (Pot Marigold)

Culture II. Cover $\frac{1}{4}$ -in. Ht. 18 to 20-in. Midsummer to frost. Grows anywhere. Stands drought and heat. Transplant to 12 to 16-in. apart.

Campfire Improved—Double 4-in. flowers, deep orange. Heavy stems. Pkt. 15c.

Orange King—Extra fine, deep orange, dark center.

Orange Shaggy—All-America 1935. Deep orange, lighter centers; Chrysanthemum-like blooms.

Mixed—

CALLIOPSIS (a) (Tickseed)

Culture IV. Cover $\frac{1}{8}$ -in. Ht. various. Blooms all summer. Bright varied combinations of color.

Golden Crown—All-America 1938. Improved rich yellow, maroon center. Pkt. 15c.

Tall, Mixed—Finest large and small-flowered varieties.

CAMPANULA (Canterbury Bells) (a)—**Culture I.** Cover $\frac{1}{8}$ -in. Ht. $1\frac{1}{2}$ ft. Blooms 5 to 6 months after seeding. Bushy plants, bell-shaped flowers, white, pink, blue and lilac.

Single—All colors.

Single and Double Mixed—Fine for late summer borders.

CANDYTUFT (Iberis) (a)—**ANNUALS:** **Culture III.** Cover $\frac{1}{4}$ -in. Ht. 6 to 12-in. Blooms in 8 weeks. Blooms all season if not allowed to seed. Will grow in dry, sandy soil.

All Flower Seeds

10c PER PACKET

except where otherwise noted

White Hyacinth—Large heads of pure white flowers.

Umbellata Mixed—Splendid range of colors.

CANDYTUFT (hp)

Culture VII. Cover $\frac{1}{4}$ -in. Ht. 8 to 10-in. Blooms in early spring and summer. Will grow in dry, sandy soil. Dwarf plants, a mass of flowers, pink, lavender or white.

CARNATION (a)—**Culture I.** Cover $\frac{1}{8}$ -in. Ht. 15-in.

Chaubaud's Giant—Double, extra large, clove-scented. Blooms 5 months after seeding. Pkt. 15c.

CASTOR BEAN (Ricinus) (a)—**Culture II.** Cover seeds $\frac{1}{2}$ -in. Broad tropical leaves and brilliant colored fruits. Grows 6 to 15 ft. Needs good soil, ample sun and moisture.

CELOSIA (Cockscomb) (a)

Culture I. Cover $\frac{1}{8}$ -in. Ht. various. Midsummer to frost.


Childsii Type (Chinese Woolflower)—Forms large heads like wool chenille balls of a bright, rich red. Ht. 2-ft.

CENTAUREA—See Bachelor's Button.

CHRYSANTHEMUM (a)—**Culture II.** Cover $\frac{1}{8}$ -in. Ht. 2-ft. Favorite flower for fall bloom. Pleasing color range. Give plenty of room.

Single Mixed—

CLARKIA (a)—**Culture IV.** Cover $\frac{1}{4}$ -in. Ht. 2-ft. Graceful, erect, bushy plants, many rosette-like flowers set along stem.



IPOMOEA, Pearly Gates

CLEOME (Spider Plant), Pink Queen (a)—**Culture II.** Cover $\frac{1}{4}$ -in. 1942 All-America. A tall, attractive border plant, free from insects and disease, with huge heads of delightful and refreshing pink, without a trace of blue or magenta. The bracts turn to white before falling. Pkt. 15c.

CLOVER STRAWFLOWER—See Everlasting Flowers.

COCKSCOMB—See Celosia.

COREOPSIS (hp)—**Culture VII.** Cover seeds $\frac{1}{4}$ -in. Ht. 2-ft. Long blooming season. Daisy-like yellow flowers with broad fluted petals. Long stems.

CORNFLOWER—See Bachelor's Button.

COSMOS (a)

Culture IV. Cover seeds $\frac{1}{4}$ -in.

Dazzler—Rich deep, velvety crimson maroon Sensation. Immense flowers, 4 to 5-in. across, with broad, heavy, fluted petals. 1943 All-America. Pkt. 25c.

Orange Flare—All-America 1935. Early flowering. Vivid orange blooms, 2 to 3-ft. plants.

Orange Ruffles—Semi-double, large, vivid blossoms. Very free blooming over long period, if old flowers are removed. 10 days earlier than Orange Flare. 3 to 4-ft. plants. Pkt. 15c.

Sensation Mixed—Largest, earliest. 4-ft. tall; flowers up to 5-in. across. All-America 1936.

Yellow Flare—All-America 1942. First yellow Cosmos ever developed. Single, 2 to 2½-in. flowers, produced in great masses on 3-ft. plants. Slightly later than Orange Flare. Pkt. 15c.

CYNOGLOSSUM (Chinese Forget-Me-Not) (a)—**Culture II.** Cover seeds $\frac{1}{8}$ -in.

Firmament—New dwarf type, with bushy plants and bright blue color. All-America 1939. Pkt. 15c.

DAHLIA (a)—**Culture I.** Cover seeds $\frac{1}{4}$ -in. Usually grown from roots, but interesting to raise from seed.

Unwin's Ideal Bedding—Miniature double and semi-double colorful flowers; dwarf bushy 18 to 24-in. plants. Long-lasting cut flower. Pkt. 15c.

DAISY, AFRICAN—See Arctotis.

DELPHINIUM (hp) ♦

Culture VII. Cover $\frac{1}{8}$ -in.

Giant Pacific Hybrids—Very large, well-shaped florets, tall, slender spikes. Pkt. 50c.

Wrexham Hybrids—Hollyhock-flowered strain. Pkt. 35c.

DIANTHUS (Pinks) (a)

Culture IV. Cover $\frac{1}{8}$ -in. Ht. 6 to 12-in. Blooms all summer.

Single Mixed—

Double Mixed—

DIDISCUS—See Blue Lace.

DIGITALIS—See Foxglove.

ESCHSCHOLTZIA (California Poppy) (a) ♦—**Culture III.** Cover $\frac{1}{8}$ -in. Ht. 1-ft. Midsummer to frost. Prefers sun. Large, silky flowers, red, gold, orange, scarlet, chestnut, pink. Lacy foliage.


EUPHORBIA—See Snow-on-the-Mountain.

EVERLASTING FLOWER (a)


(Globe Amaranth, Gomphrena Everlasting or Clover Strawflower)—**Culture II.** Cover $\frac{1}{8}$ -in. Ht. various. Blooms all summer. Retains natural colors when cut in bud and dried. A true Bachelor's Button that may be dried.

Strawflower (Helichrysum)—Large double flowers. Assorted.

FOUR O'CLOCK (a) ♦—**Culture III.** Cover $\frac{1}{4}$ -in. Ht. 2-ft. Thrives under



CAMPANULA



ESCHSCHOLTZIA

any conditions. Sweet scented. Red, yellow, rose, white, lilac blossoms open in mid-afternoon.

FOXGLOVE (Digitalis) (b) ♦—**Culture VII.** Cover seeds $\frac{1}{16}$ -in. Height 3 to 7-ft. Hardy Biennials for tall borders. Tall spikes of bell-shaped blossoms. All colors.

GAILLARDIA (a) (Blanket Flower)

ANNUALS: **Culture II.** Cover $\frac{1}{8}$ -in. Ht. 1½-ft. Blooms all summer. Needs full sunshine.

Single Mixed—

Double Mixed—

GAILLARDIA (hp)

PERENNIALS: **Culture VII.** Cover $\frac{1}{8}$ -in. Ht. 2-ft. Showy blooms in orange, yellow, red and maroon. Withstands drought, heat.

Perennial Monarch Mixed—Large range of colors.

GEUM (hp)—**Culture VII.** Cover $\frac{1}{8}$ -in. Ht. 2-ft. Flowers like tiny double roses, long stems, prized for cutting. Pkt. 15c.

GILLIFLOWER—See Stocks.

GLOBE AMARANTH—See Everlasting Flowers.

GODETIA (Satin Flowers) (a) ♦—**Culture II.** Cover $\frac{1}{8}$ -in. Ht. 1-ft. Bushy plants with masses of Azalea-like flowers. Thrives best in moist soil and half-shade.


Double Mixed—Long spikes with pom-pom-like blossoms.

GOARDS, ORNAMENTAL (a)—**Culture II.** Cover seeds $\frac{1}{2}$ -in. Useful climbers, large foliage; singularly shaped. Last years when fully ripened and cured. Mixed.

GYPSOPHILA (a) (Baby's Breath)

ANNUALS: **Culture II.** Cover $\frac{1}{8}$ -in. Ht. 1½-ft. Blooms 6 to 7 weeks. Sow two weeks apart for continuous bloom.


White—A pure strain.



DIANTHUS
Single



PHLOX, Rosy Morn



DAHLIA UNWINS
Dwarf Hybrids


GARDEN GUIDE POSTS



A vegetable garden only
25 x 50 feet will keep a
hungry family of five in
fresh vegetables all through
the season — with plenty
left over for canning.

flower, broccoli and kohlrabi, allow thermometer to go below 75 degrees.


On sunny days ventilate thoroughly, but even on cloudy days a crack of air may be needed to get rid of ammonia odor. Cover glass at night with blankets, old rugs or hotbed mats to conserve heat.



Basement Window Greenhouse


The frame for a basement window greenhouse is made the same as for a manure hotbed, but the back is omitted and no pit is dug. The open back fits against the basement window. The open window both ventilates and heats, so that in most instances no further ventilation will be needed. On very cold nights, a 100 watt electric bulb will keep up heat.

Electric hotbed cables are usually installed over a bed of sand six inches thick, and covered with $\frac{1}{2}$ " mesh hardware cloth. Electric lights (two 100 watt lamps for each 3 x 6 sash) can be used and controlled by a chicken brooder thermostat.



Cold Frames

A cold frame is the same as a hotbed without manure or other source of artificial heat. It depends upon the sun for heat, and must be covered with hotbed mats to conserve this heat at night. It can be used only after the weather has warmed up somewhat, but before the danger from late frosts is past. Lettuce and radishes planted in the cold frame a month before frost is expected can be grown until late in the year if the glass is kept covered at night.



If manure is not available, dried leaves or chopped straw may be used by adding a handful of ammonium nitrate or ammonium sulfate to each bushel of leaves. This will not, however, generate as high a heat as manure.

As soon as frame is in place, water well, insert thermometer in soil and cover with sash, airing daily until thermometer drops below 85 degrees. This is time to plant tomatoes and peppers. For cabbage, cauli-




LARKSPUR, Giant Imperial Lilac King



HELIANTHUS, Sun Gold



ZINNIA, Dahlia-Flowered Exquisite



MARIGOLD, Sunkist

GYPSOPHILA (hp)

PERENNIALS: Culture VII. Cover $\frac{1}{8}$ -in.

Perennial White—Tiny flowers on slender stems, producing a mist-like effect.

HELIANTHUS (Sunflower) (a)—Culture II. Cover $\frac{1}{4}$ -in.

Sun Gold—New! Huge 100% double flowers, free blooming, 4 to 5-ft. tall. Pkt. 15c.

HELICHRYSUM — See Everlasting Flowers.

HEUCHERA (Coral Bells) (hp) — Bright coral red flowers on wiry stalks. Charming for low border or rock garden. Pkt. 15c.

HOLLYHOCK (hp)

Culture VII. Cover seeds $\frac{1}{4}$ -in. Ht. 5 to 8-ft. Succeed best in deep, rich, well-drained soil. Like warm, sunny place.

Double Mixed —

KOCHIA (Burning Bush or Fire Bush) (a) — Culture II. Cover $\frac{1}{8}$ -in. Ht. 3-ft. Close, compact, bright green pyramidal bush, moss-like foliage that turns brilliant red in fall. Fine hedge plant.

LADY SLIPPER—See Balsam.

LARKSPUR (a)

Culture IV. Cover $\frac{1}{8}$ -in. Ht. 2 to 4-ft. Blooms all summer. Clustered flowers on tall spikes, feathery foliage. Keep faded blooms cut.

Blue Spire, Giant Imperial—Most desirable blue. Pkt. 15c.

White King, Giant Imperial—Large, double, pure white. Pkt. 15c.



GAILLARDIA

NICOTIANA

Giant Imperial Mixed—Pkt. 15c.

Pink Perfection—2-in. florets, cutting spike 24-in. or more. Early flowering. Pkt. 15c.

LINUM (a) ANNUALS: Culture III. Cover $\frac{1}{8}$ -in. Ht. 15-in. Need sun.

Scarlet Flax—Brilliant glossy-red flowers on delicate stems. Life of individual flower short, but blooming season can be prolonged by successive sowings.

LINUM (hp) ♦ —PERENNIAL: Culture VII. Cover $\frac{1}{8}$ -in. Need sun. Ht. 15-in.

Perennial Blue—Light blue, upright flowers in great profusion all summer.

LOBELIA, Crystal Palace (a)—Culture V. Cover $\frac{1}{16}$ -in. Ht. 4-in. Small, round, compact plants covered with deep blue flowers, dark foliage.

LUPINES (a) ♦—Culture III. Cover seeds $\frac{1}{2}$ -in. Midsummer. Prefers light shade. Long spikes of large dark blue, sky blue, rose, or white flowers. Mixed.

LUPINS (hp)

Russell's Choice Mixed—Wonderful border flower of many lovely colors. Gold Medal winner, 1937. Pkt. 25c.

MARIGOLD (a)

Culture II. Cover $\frac{1}{8}$ -in. Ht. various. Blooms early and late.

Burpee's Gold — Extremely large, sweet-scented, double. Odorless foliage. 2 $\frac{1}{2}$ -ft. Pkt. 15c.

Butterball, Dwarf French Double—All-America 1942. Compact, 6 to 10-in. plants; very early. 1 $\frac{1}{2}$ -in. blooms. Pkt. 15c.

Crown of Gold, Odorless—All-America 1937. Short, quilled center petals, with flat, deep orange guard petals. Branching plant, 2 $\frac{1}{2}$ to 3-ft. Midseason.

Dwarf Royal Scot — Mahogany red flowers with gold stripes. Plants, 24 to 30-in. Pkt. 15c.

Guinea Gold—Magnificent golden flowers with carnation-like petals. Plants 2 to 2 $\frac{1}{2}$ -ft.

Harmony Type, French Dwarf Double Hybrids—Flowers 1 $\frac{3}{4}$ -in., all double. No strong Marigold odor. Plants 12 to 14-in. Combinations of yellow, orange, blotched and striped combinations of orange or yellow with maroon. Earliest of all. Pkt. 15c.

Harmony Type, French Tall Single, Ferdinand — A tall French Marigold with brown rays and golden tufts. Pkt. 15c.

Spry—1941 All-America. Extra dwarf double French type; about 9-in. tall; 1 $\frac{1}{2}$ -in. flowers, clear yellow crested center, mahogany-red guard petals. Blooms in 9 weeks. Pkt. 15c.

Sunkist — 1943 All-America. Dwarf, compact, very early. Continuous blooming, 6 to 10-in. mound-shaped plants. Flowers to 1 $\frac{1}{2}$ -in. across. Pkt. 25c.

Yellow Supreme—Sweet-scented, yellow, 3-in. blossoms.

Yellowstone, Mission Giant — All-America 1942. Late blooming. Pkt. 20c.

MIGNONETTE, Sweet Scented (a)—Culture I. Cover $\frac{1}{16}$ -in. Ht. 15-in. Dwarf, with many small spikes of fragrant blossoms. Long blooming.

MOON FLOWER (a)—Culture VIII. Cover seeds $\frac{1}{4}$ -in. Plant in a sunny location. Quick-growing vine with dense green foliage and pure white flowers, 5 to 6-in. across, that open in evening and on cloudy days. Fragrant.

MORNING GLORY (a)

(Ipomoea)

Culture III. Cover $\frac{1}{4}$ -in.

Heavenly Blue—Vigorous, quick growing, with extremely large blue flowers. Long blooming. Pkt. 15c.

Scarlett O'Hara — All-America 1939. Rich wine-red or deep rosy crimson blossoms about 4-in. across. Pkt. 15c.

Pearly Gates—1942 All-America. Unusually large white flowers, luxurious foliage. Plant with Heavenly Blue and Scarlett O'Hara for red, white and blue effect. Pkt. 15c.

Tall Mixed — A very large-flowered strain with many lovely colors.

MOSS ROSE—See Portulaca.

MOURNING BRIDE—See Scabiosa.

NASTURTIUM (a)

Culture III. Cover $\frac{1}{2}$ -in. Prefers sunny location. Thrives in poor soil.

Dbl. Glorious Gleam Hybrids—Sweet scented. Long stems. Compact habit. Many colors.

Golden Gleam—Double and semi-double fragrant gold blossoms, semi-tall plants.

Golden Globe (Dwarf Dbl.) — Color like Golden Gleam. Dwarf.

Scarlet Gleam—Fiery red.

Single Tall or Climbing—May be used for covering fences or trailing over rockwork. A wide range of colors.

NEMOPHILA (Baby Blue Eyes) (a) —Culture II. Ht. 6-in. Dwarf, spreading plants; sky blue flowers, lighter toward center.

NICOTIANA (a) ♦—Culture I. Cover $\frac{1}{16}$ -in. Ht. 2 $\frac{1}{2}$ to 3-ft. Long tubed, petunia-like blossoms pour rich, spicy fragrance over the garden morning and evening.

PANSY (a) ♦ — Culture VII. Cover $\frac{1}{8}$ -in. Ht. 6-in.

Swiss Giants — Long-stemmed, especially for cutting. Pkt. 50c.

PETUNIA (a)

Culture I. Cover $\frac{1}{16}$ -in. Ht. various. Blooms summer and fall.

Burgundy—A rich wine red with contrasting white throat. Single plain edged.

Dainty Lady—All-America 1936. Large, single fringed, yellow, ringed gold and cream; compact plants. Pkt. 35c.

Flaming Velvet — Dark red, single, plain edged. Pkt. 15c.

Fluffy Ruffles—A delightful mixture of delicate light shades. Finely fringed and very heavily ruffled so that they appear double. Finest choice for home gardens.

Giants of California, Mixed—A well balanced mixture of large fringed and ruffled flowers of light and dark shades. Suitable for window box or potting.

Glamour—1942 All-America. Huge 5 to 6-in. blooms; appear when plant only 5 to 6-in. tall. First appear double, but as they mature, petals fall back, exposing throat. Plants 1 $\frac{1}{2}$ to 2-ft. Pkt. 25c.

Rose of Heaven — Rose, with inconspicuous light throat. Single flowers. dwarf plants.

Rosy Morn—Medium sized rose-pink flowers, white throat. Plants, 1-ft. Pkt. 15c.

Single Fringed, Mixed — Excellent mixture of rich bright colors. Medium size and erect growing.

Single, Plain Edged, Mixed—Special mixture, light and dark shades.

Single, Plain Edged, White — Pure white, dwarf, compact.

Theodosia—Flowers 3 to 3 $\frac{1}{2}$ -in., rose, with gold throat; single fringed. Flowers when plant very small. Pkt. 25c.

All Flower Seeds

10c PER PACKET

except where otherwise noted

GARDEN GUIDE POSTS

PREPARING THE GARDEN SOIL

A good gardener can be told by the way he treats the soil. No gardener worthy of the name works it when it is so wet that it will pack. Abuse of the soil may ruin it for the entire season. Use the mud pie test already described before digging or plowing.

If your soil is heavy clay, lime added before plowing will help break it up. Even more valuable is decayed vegetable matter of some sort—peat moss, humus, old manure or leaf mould.

Do not apply fresh manure to green vegetation in spring because, as this decays, it will form a gas that will hurt roots.
If necessary to make a garden on sod, skin off the sod and add it to the compost heap. If the garden is fall-plowed, however, sod can be turned under and will decay fairly well if lime and fertilizer are applied at the same time.

To Dig or Plow

Many a gardener has wasted precious time waiting for a plowman to turn over a garden that could easily be turned over with a spade and would be in better condition than if packed by a heavy tractor. Hand digging, if properly done, is better for a garden than tractor plowing. Larger areas, of course, must be worked by machinery, but if possible, do the final finishing by hand.

The trick of hand spading is to take your time. A good sized garden can be turned over in about a week by working at it an hour a day. When you dig, don't cheat the spade. Drive it straight down, lift the entire clod and turn it completely upside down, hit it with the back of a spade, and if the soil is in good condition it will crumble.



To be sure that each clod is actually turned over, remove a trench along the beginning of the garden just one spade wide. Then as each spadeful is lifted, turn it over into this trench.



Vegetables? Of course! But this year plant plenty of flowers, too!

The soil removed from the first row is wheeled around to where you plan to finish, and is used to fill the last trench or spit.

Trenching (the method used by these) does not pay for a temporary garden, but should be practiced by home owners who plan to work the same plot year after year. In this method a trench is dug two full spades deep.

Manure or compost is spread over the top of the entire garden and also at the bottom of each trench as dug. The second row of soil is thrown into the bottom of the trench and the layer just below it is thrown on top of it. In this way the entire garden is turned upside down and the ground is loose and friable to a depth of nearly two feet.

Because considerable subsoil is mixed with the upper layer, an extra application of fertilizer to the garden after digging is good practice.

From the second year on, the soil should get better and better, even though it is only turned over one spade deep each year.

How Much Fertilizer? What Kind?

Well rotted manure and compost are important to the production of good crops and they do feed plants, but it must be remembered that they do not contain the type of nitrogen that is needed early in the season.

If 30 lbs. of a good mixed fertilizer are applied to every 1,000 feet of garden before digging, and an equal amount applied between the rows about half-way through the season, all of the plant food needs of the average garden can be met.

Always "water in" the summer application to prevent burning.

PLANTING THE SEED

Don't try to make rows without a line or straight edge. Place a stake at each end of the row and stretch a line tightly between them. Make the furrow for the seed with a hoe.

The old theory that seed should be planted deep enough to come in contact with soil moisture still holds, but in dry weather the soil should be watered thoroughly before planting so the moisture will be right at the surface. If the row is then covered with a board until the seed germinates, it will not be necessary to bury the seed even though the weather is dry.

Fine seed (onion, lettuce, cabbage, etc.) should not be planted more than $\frac{1}{2}$ " deep. Medium seeds (okra, spinach, cucumber) 1" deep. In heavy soils, beans, squash, onion sets and peas need not be covered more than 1" but the soil must be moist. In lighter soils, 2" is deep enough.



STRAWFLOWER, Everlasting



CALLIOPSIS
Tickseed



GODETIA, Mixed



ZINNIA
Harmony
Type

GARDEN GUIDE POSTS

HOW?
WHEN?
WHERE?
WHY?



CALENDULA, Orange Shaggy



MARIGOLD, Harmony
Type, French
Tall Single



MARIGOLD, Guinea Gold

(Below) GAILLARDIA



SEED PLANTING (Cont'd)

One of the common errors in planting squash, melon, cucumber, etc., is to assume that a hill really is a hill. Actually, a hill is a hole, since these vine crops are planted in depressions about 2" below the surrounding surface to catch water. Incidentally, all vine crops love humus or well-rotted manure. If they can be planted in depressions filled with soil in the top of a compost heap, they will produce extravagantly.

Good seed germinates freely, so it is a mistake to plant it too thickly. This is especially true if it is treated before planting with ROOTONE, which speeds up germination and root growth.

How to plant fine seed

In planting fine seed, tear off one corner of the packet and, if Rootone is used, drop in a pinch and shake. The seed can be tapped out of the torn corner of the packet one at a time. Fine seed is often mixed with dry sand so that it will spread out better, but this will only work if seed is not more than twice the size of the sand grains.

Large seeds should be poured out of the packet into a cup or bowl so that they can be dropped individually into the furrow. Beans and peas should be treated with a legume inoculation before planting, even if this was done last year. Improved legume cultures are constantly being propagated in the laboratories and give superior results to natural inoculation.

TIME OF SOWING

The danger from frost or cold and wet weather varies so much that exact dates are impossible to give. However, the following natural calendar will tell when various crops are to be planted.

FULLY HARDY CROPS: Sow as soon as the ground can be worked in spring, or when the farmer plants oats. In some years crocuses and snowdrops may be in bloom at this time, but if the ground is ready sooner, plant anyway. These crops include lettuce, cabbage, broccoli, carrots, cauliflower, kale, kohlrabi, parsnips, radishes, rutabagas, spinach and turnips. Also round-seeded peas.

HARDY CROPS: When the daffodils and earliest tulips bloom, or when maple trees are in full flower, plant these: Beets, wrinkled peas, Swiss chard, and onion sets.

TENDER CROPS: When the Liliacs, Darwin Tulips and Apple Blossoms are in full bloom, plant these: Bush beans, sweet corn, pumpkins, and squash are included.

TROPICAL CROPS: When the petals on apple blossoms will have fallen and the tall late irises are in full bloom plant: Okra, Lima beans, Pole beans, muskmelons and cucumbers, and set out plants of tomatoes, peppers and eggplant. These can't stand frost and should not be planted until the weather is warm and settled.



TRANSPLANTING

Seedlings can be transplanted any time after the second pair of leaves has formed, provided the weather is suitable.


The effects of "hardening off" seedlings by exposing them to cool air is not very effective; even a fully hardened cabbage plant will lose its frost resistance if subjected to warm weather for four or five days. Therefore, even in the case of crops that are seeded very early, seedling plants should not be set outdoors until the weather is warm and settled.

Cabbage, broccoli and cauliflower can go out when sweet corn is planted, and tomatoes and peppers after apple blossom petals fall.

Wet the soil around seedlings thoroughly before planting out. Select a cloudy day so the direct sun won't wilt the plants. Dig with a ball of soil that contains as many fine hair roots as possible.

Watering the seedlings with a solution of TRANSPLANTONE when transplanting will restore the hormones lost when the root tips are cut in digging. This plant hormone prevents wilting and means quicker recovery from the shock of transplanting.

In working between rows when transplanting, lay a board over the soil so it won't pack. Roots grow better in loose soil.



Right Wrong Wrong

When seedlings are to be transplanted without a ball of soil, long taps roots should be shortened, and the whole root system placed and "firmed in" so that it is well spread out, not bunched. Soil in which roots are placed must be finely pulverized.



Use a wide board when transplanting, to prevent packing soil around roots. (See sketch.)



SCABIOSA, Blue Moon

PHLOX (a) ♦

Culture II. Cover $\frac{1}{8}$ -in. Ht. various. Prefers sun. Thrives in almost any soil. Art Shades—A wonderful range of soft shades. Light and deep salmon with cream eye, pale violet and soft rose with white eye. A beautiful assortment.

Rosy Morn—A bright color combination of deep rose pink with a white eye. A free bloomer.

Salmon Glory—A deep salmon with cream eye.

POPPY (a)

ANNUALS: Cover $\frac{1}{8}$ -in. Ht. various. Grow freely in almost any soil. Prefer sun.

California—See *Eschscholtzia*.

Iceland Mixed (hp)—Culture VII. Blooms first year. Excellent color balance. Immense blooms and very long stems. 2½ to 3-ft. Pkt. 15c.

Shirley Mixed—Ruffled edges, large blooms.

PORTULACA (Moss Rose) (a)—Culture III. Cover $\frac{1}{8}$ -in. Ht. 5-in. Thrive in sunny location and light soil. Fine for mass planting, edgings.

Double Mixed—Pkt. 15c.

Single Mixed—

POT MARIGOLD—See *Calendula*.

RICINUS—See *Castor Bean*.

SALPIGLOSSIS (Painted Tongue) (a)—Culture II. Cover $\frac{1}{16}$ -in. Ht. 2½-ft. Blooms midsummer to frost. Large, lily-like flowers, velvet textured, in many bright colors.

SALVIA (Flowering Sage) (hp)—Lavender-blue flowers on plants about 3-ft. high. This perennial will bloom first year if planted in spring and lasts until the first frost. Pkt. 15c.

SATIN FLOWER—See *Godetia*.

SCABIOSA (a)

(Mourning Bride)

Culture II. Cover $\frac{1}{8}$ -in. Ht. 2½-ft. Early summer to frost.

Blue Moon, Imperial Giants—Double flowers, resembling old-fashioned beehive. Lavender blue. All-America 1939. Pkt. 25c.

Rosette—Very large; rose, flushed with salmon. Pkt. 15c.

Mixed—

SCARLET FLAX—See *Linum*.

SCHIZANTHUS (Butterfly-flower or Poor-Man's Orchid) (a)—Culture II. Cover $\frac{1}{8}$ -in. Ht. 2-ft. Flowers like tiny butterflies in pastel shades, with dainty markings. Finely lacinated foliage.

SHASTA DAISY, Alaska (hp)—Culture VII. Cover $\frac{1}{8}$ -in. Ht. 2-ft. White single blossoms, 4 to 5-in. across, with small yellow centers, long stems.

SNOW-ON-THE-MOUNTAIN (Euphorbia) (a)—Culture IV. Cover $\frac{1}{8}$ -in. Ht. 2-ft. Foliage beautifully edged with white. Prefers sun.

STATICE (Sea Lavender) (a)—Culture I. Cover $\frac{1}{8}$ -in. Ht. 1½-ft. Seed germinates very slowly. Masses of dainty flowers in large panicles, giving a mist-like effect. When dried, flowers last for months.

STRAWFLOWER—See *Everlasting*.

STOCKS (a) ♦

Culture I. Cover $\frac{1}{8}$ -in. Ht. 2 ft. Bloom all summer. Like well-enriched soil, plenty of moisture, some shade in hot months.

Giant Imperial—Large flowers, long stems. Very early; many doubles. Pkt. 25c.

Ten Week Stocks—Double rosette-like blooms, fine stems.

Beauty Mixed—

SUNFLOWER—See *Helianthus*.



ZINNIA
Scabious Flowered

ZINNIA
Lilliput

SWEET PEAS (a)

Sow very early. Dig trench 1-ft. wide and deep. Place 6-in. rotted manure or peat moss mixed with manure, on bottom, tread down firmly. Fill with good garden soil. Sow one row of Sweet Peas $\frac{3}{4}$ -in. apart. When plants are 3 to 4-in. high, thin to 6-in. apart and provide supports. Need plenty of water; and mulch of peat moss when warm weather approaches. Pick flowers daily.

Early Flowering Spencer—Individual colors and mixed. Can be grown either inside or out of doors.

Late or Summer Flowering Spencer—Individual colors, mixed. Extremely hardy, productive, and more heat resistant than other varieties.

Ruffled Spencer—Individual colors, mixed. Contains many varieties in a perfect blend of color.

TICKSEED—See *Calliopsis*.

TITHONIA, Fireball (a)—Culture I. Cover $\frac{1}{8}$ -in. Ht. 6 to 8-ft. Blooms in late summer. Dazzling scarlet, large single flowers. Plant very tall.

TRITOMA (Red Hot Poker) (hp)—Striking red spires of bloom from August on. About 3 ft. tall. Continuous blooming. Newly popular.

VERBENA (a)

Culture I. Cover seeds $\frac{1}{8}$ -in. Ht. 15-in. Dwarf plants with very large heads.

Mammoth White—

Scarlet—

Shades of Blue—

Mixed—

WILD FLOWER MIXTURE—Splendid mixture of old fashioned garden flowers, various kinds and colors, for interesting border planting.

ZINNIA (a)

Culture III. Cover $\frac{1}{8}$ -in.

Black Ruby—1941 All-America. Deep velvety maroon. Lilliput or Pompon zinnia of merit. Distinct color for this class; 1¼ in. flowers, long stems. Plants 18 in. tall. Effective used with lighter colors. Pkt. 15c.

Crown of Gold—Giant flowers, petals deep golden yellow at base, with individual colors at tip. Plants 2-ft. Pkt. 15c.

Dahlia Flowered—6-in. blooms. Plants 2½ to 3-ft. Most satisfactory Zinnia. Yellow, red, pink, orange. Pkt. 15c.

Fantasy—Great masses of shaggy ray-like twisted petals; 2-ft. plants. **Melody** (orchid-lavender), pkt. 15c; **Star Dust** (clear golden yellow), pkt. 15c; **Wildfire** (rich scarlet), pkt. 20c; **White Light** (pure white), pkt. 15c; **Mixed**, pkt. 10c.

Giants of California—Huge blossoms, on 3-ft. plants. Variety of colors. **Mixed**.

Harmony Type, Mixed—Plants 3 ft. bearing numerous medium sized flowers. Fine mixture contains bright shades such as crimson, orange, yellow, pink and white.

Lilliput—Straight-stemmed midget pompons. Red, yellow, pink, white, orange and mixed.

Lilliput Pastel Mixture—Plants 12 to 18-in. Pompons 1 to 1½-in. Variegated pastel colors.

Linearis—Single, golden-orange flowers, lemon yellow stripe, brown centers. Dwarf 1-ft. plants. Pkt. 15c.

Mexicana—Miniature yellow, orange and mahogany flowers, frequently variegated. Single and double, on 1-ft. plants.

Navajo Mixed—Bi-colored. Medium sized flowers; 2½ to 3-ft. plants. All-America 1938. Pkt. 15c.

Pumila Sunshine Tints—Symmetrical 2½ to 3-in. flowers, in luscious autumn tints and pastel shades. Very free flowering. Long stems, fine for cutting. Pkt. 15c.

Scabious Flowered—Large semi-globular crown, surrounded by fringe of petals. Plants 2½ to 3-ft. Pkt. 15c.

Super Crown-O'Gold, Pastel Tints—Flowers 5 to 5½-in. Wide range of pastel shades. Pkt. 25c.

Will Rogers—A striking new red, never before available in Zinnias. Very robust plants, literally loaded with mammoth blooms, 6 inches across, and often 4 inches in depth, resembling the Show Dahlia.

All Flower Seeds
10c PER PACKET

except where otherwise noted

Planting Guide... FOR THE FLOWER GARDEN

KEY—For classification of flowers by purpose, location, etc., see the second column below. The meaning of the key letters in this column is as follows:

A—For dry, hot locations.
B—For a succession of blooms.
C—For shady locations.
D—For poor soil.
E—For edging.
F—For the rock garden.

G—For their foliage.
H—Everlasting for winter bouquets.
I—For their fragrance.
J—For porch or window boxes.
K—Climbing vines.
L—For backgrounds.

	Height (inches)	Suitable for	Average Days to Germinate	Approx. Time of Bloom	Shape of Plant						
Ageratum	4-12	E F J	5-10	10 Wks.	bushy	Godetia	12-24	C D	15-20	2 Mos.	erect
Alyssum	2-12	B C D E F I J	5-10	6 Wks.	spreading	Gypsophila	18-36	B H	15-20	8-10 Wks.	branching
Antirrhinum	12-60	C I L	20-25	3-4 Mos.	bushy	Helichrysum	24-36	H	5-10	3 Mos.	slender
Aquilegia	18-36	C E F	15	3 Mos.	branchy	Hollyhock	48-84	L	5-20	10 Wks.	bushy
Balsam	14-20	A E	10-12	6 Mos.	erect	Kochia	30-36	A B	15-18	Foliage	tall
Calendula	12-36	D	10-14	12-14 Wks.	bushy	Larkspur, (A)	30-72	L	15-20	10-12 Wks.	vine
Campanula	18-24	F	8-15	10-12 Wks.	bushy	Lathyrus	60-72	I K	25-40	2-3 Mos.	vine
Candytuft	6-18	B F I	5-10	8-10 Wks.	bushy	Lobelia	4-8	E J	10	10-12 Wks.	bushy
Centaurea	18-36	A B C D E I J	5-20	3-4 Mos.	erect	Marigold	8-36	B J	5-8	6-8 Wks.	bushy
Chrysanthemum	24-36	A E	5-20	8-10 Mos.	bushy	Mignonette	10-12	B I	3-4 Mos.	erect	
Celosia	12-36	D E H	20-25	10 Wks.	bushy	Moon Flower	12 ft. plus	I K	5-8	4-5 Mos.	vine
Cosmos	36-60	A	5-15	10-12 Wks.	bushy	Morning Glory	12 ft. plus	A K	5-8	3-4 Mos.	vine
Coreopsis	18-36	A B D	10-20	4-5 Mos.	bushy	Nasturtium	12-36	D E F K	8-15	2-3 Mos.	bushy vine
Dahlia, Dwarf	14	F	10	3-4 Mos.	bushy	Nigella	10-18	B F	10-15	10-12 Wks.	bushy
Delphinium	36-72	L	15-21	3-4 Mos.	tall	Nicotiana	30-48	I	20-25	10-12 Wks.	branchy
Digitalis	30-48	F	10-20	3-4 Mos.	erect	Pansy	4-8	C E F	10-15	10-12 Wks.	clump
Dianthus	4-15	F I	5-10	12 Wks.	bushy	Petunia	10-24	D E F J	18-20	10-12 Wks.	bushy
Eschscholtzia	10-12	C D F	5-12	12-15 Wks.	bushy	Phlox	10-18	A B C F I J	20-25	8-10 Wks.	bushy
Four O'Clock	24-30	A D	5-10	3-4 Mos.	bushy	Poppy	18-36	B	20	12-15 Wks.	erect
Gaillardia	18-30	D	15-20	2-3 Mos.	bushy	Portulaca	4-6	A D E F J	18-20	6 Wks.	spready
						Salvia	12-42	A	15-25	3-4 Mos.	bushy
						Scabiosa	18-30		18-20	2-3 Mos.	bushy
						Stocks	12-24		5-15	12-14 Wks.	bushy
						Schizanthus	18-36	C E	20-25	6-8 Wks.	bushy
						Sweet Pea	36 or more	K	15-20	10-12 Wks.	vine
						Sweet William	12-24	F	10	12 Wks.	bushy
						Verbena	6-12	E F J	8-10	12 Wks.	spready
						Wallflower	12-18	I	5-12	2-3 Mos.	bushy
						Zinnia	12-36	A F	5-10	6-8 Wks.	bushy

The 8 CULTURES... SEE CULTURE NUMBERS IN OUR FLOWER LISTING

CULTURE I

Seeds give best returns if sown in flats in early spring, but good results may be obtained by sowing outdoors after all danger of frost is past. Sow seed at depth indicated under flower descriptions. Water thoroughly with fine spray and place in warm room; watch flats carefully and never let soil become dry. When seedlings appear, remove flats to sunny window. Transplant to other flats as soon as they are large enough to handle, keeping them 2" to 2½" apart both ways. Expose gradually to outside conditions and, when sufficient growth has been made, plant in permanent outside locations. For sowing outdoors, follow instructions given in Culture II.

CULTURE II

May be sown in outside seed-beds and later transplanted, or directly into their permanent location. Spade seed-bed deeply, mix well-rotted manure, or peat moss or humus and a commercial fertilizer, well into the soil; spread an additional half inch of peat moss or humus on seed-bed, and rake well into top surface. Sow seeds at depth indicated, and water thoroughly with a fine spray. When large enough, transplant or thin out. If exceptionally early flowering is desired, follow instructions under Culture I.

CULTURE III

Seeds do not take kindly to transplanting and should be sown outdoors, as soon as danger of frost is over, in beds they are to occupy. Prepare seed-bed same as Culture II. Level off and sow seeds thinly at depth indicated; water thoroughly with a fine spray. After seedlings appear, thin out.

CULTURE IV

Seeds under this culture, being quite hardy, may be sown outside in very early spring, as frosts do not affect their growth to any extent. The seed-bed should be spaded deeply, and fertilizer mixed thoroughly with the subsoil. Plant at depth indicated, and water thoroughly with a fine spray. When large enough to handle, thin out and transplant. For very early plants, sow in fall, leaving plants to winter in their beds.

CULTURE V

Sow in flats in early spring if for the garden; or during the summer if you wish to cultivate them in pots for house decoration. Sow seeds to depth indicated, and water thoroughly with a fine spray. Place in a warm room. As soon as seedlings appear, remove to a sunny window. When large enough to handle, transplant into other flats to stand 2" apart in rows. Transplant outdoors when all danger of frost is past. If for house decoration place in larger pots.

CULTURE VI

Plants are purely conservatory or hot-house plants. Follow instructions for preparation of seed-flat above and sow seeds preferably during spring months at depth indicated. Water thoroughly with a fine spray, and place in warm room. As soon as seedlings appear, remove to sunny window. When large enough to handle, transplant into other flats to stand 2" apart. Later pot up in 2" or 2½" pots and change to larger pots as often as needed.

CULTURE VII

May be sown in early summer in flats or coldframes at the depth indicated. Water thoroughly with a fine spray. Place in a

warm room and as soon as seedlings appear, remove to a sunny window. When large enough to handle, transplant into other flats, spacing seedlings 2" to 2½" either way. Later, when they begin to crowd, transplant them a second time. In fall, if well developed, they may be placed in their permanent locations where they are to blossom the following year, or left in the seed-beds for early spring transplanting. Note: Only a few varieties under this culture will produce flowers the first year, and all will benefit by winter mulch or protection at least the first season. Seed may be sown in the open, after danger of frost is past, in a seed-bed in which the top soil is finely pulverized and mixed with peat moss or humus. Raise seed-bed several inches above the rest of the garden to prevent seeds being washed out. When plants are sufficiently large, place them in their permanent locations.

CULTURE VIII

Best sown indoors in early spring directly into small pots. Fill the pots with a mixture of good garden soil, composed of two parts soil and one part humus, to within a half inch of the edge of the pot. Firm the soil and place one or two seeds at the depth indicated. Take an ordinary seed-flat, fill with moist peat moss, and plunge the pots into this up to within a half inch of the edge. Water pots thoroughly. Place in a warm room and as soon as seedlings appear, remove to a sunny window, gradually exposing them to outside conditions. When all danger of frost is over remove from pots without destroying the root-ball and plant in their respective places outdoors. If advanced blooming is not desired, seeds may be sown outside as soon as warm weather sets in.

GARDEN GUIDE POSTS



WATERING

Except for transplants, try to avoid watering the garden until it really needs water. When you do water, soak, don't sprinkle. Do a thorough job. The hose or sprinkler should be allowed to run on one spot until water stands on the surface in a pool.

After watering, the soil should be moist to a depth of five to six inches. Check this with a spade to be sure.

Watering tomatoes calls for special care. Cracks and blossom "end rot" (the free or blossom end turning black) are both caused by fluctuating water supply. Never let tomatoes be dry one week and wet the next. Tomatoes must be kept either moist constantly or be grown in dry soil and not watered at all.



Never apply water with such force that the top soil is washed away. Use a canvas hose which allows water to ooze out through the pores, or remove the nozzle of the regular hose and let the stream run out on a board. Overhead sprinklers waste some water, but do prevent washing. Watering in the evening or early morning means that the moisture sinks in and is absorbed by the soil before the sun can evaporate it.

CULTIVATION AND WEEDING

1. "Destroy them young" should be the first principle in controlling weeds. Weeds rob the soil of moisture and food. Worst of all, if allowed to grow, they shade the crops that require six hours of full sunshine to grow normally. Plan to spend half an hour a day hoeing weeds and the average garden can be kept up without difficulty.

2. Keep the soil from caking or cracking because when rain comes, it cannot penetrate the hard surface and may run off and be lost. A prong cultivator will leave the surface loose and ready to absorb rain.

3. Air is vital to plant growth: stir the soil as soon as the ground is dry enough after a rain or after watering. This also saves moisture. Use the hoe and save the hose.

4. When cultivating stay as far away from the plants as you can to avoid packing the soil around the roots.

MULCHING

A mulch of straw, leaves or lawn clippings will keep the surface moist and conserve water. It reduces soil

temperatures and prevents root injury from heat. It will also keep tomatoes, pepper and eggplant clean and will prevent soil insects from attacking melons.

However, if summer rains are light, the mulch can do harm, because the few drops that do fall cannot penetrate to the soil. In this case, pull away the mulch from around the plant stem and soak the roots thoroughly with the hose.

A mulch is particularly beneficial to potatoes because tuber formation stops in hot soils. As much as 12" of straw can be used over potato roots.

HARVESTING

The big advantage the home gardener has over the commercial grower is that he can pick his vegetables one minute and have them in the kitchen the next. All vegetables are better if left on the plant until ready to serve. Sweet corn in particular loses sweetness every minute it is off the stalk.

Pick only what vegetables you can use right away and not until you are ready to use them.

Most vegetables, particularly summer squash, sweet corn, cucumbers, carrots, beets, peas and string beans, should be used much younger than usually harvested by the commercial grower. Learn the "feel" of an ear of corn that is ready to pick. The tip of the ear will feel full and the silks should be turning brown.

Muskmelons do not get sweeter after picking. For full flavor leave them on the vine until a crack appears all around the point of attachment between the stem and melon. The free end should feel definitely soft when the melon is ready.

Avoid harvesting beans while the plants are wet with dew or rain, as moisture spreads disease.

HAVE A COMPOST PILE

To assure having sufficient humus for your garden, start a compost pile. In a corner of your garden (perhaps behind a screen of bushes) dig a trench about 1 foot deep, 2 or 3 feet wide, and 3 or 4 feet long. Make frame extending above ground level.

Humus is decomposed vegetable or organic matter. In building a compost pile the idea is to accumulate such matter in a way that will encourage decomposition and increase its plant food value.

After you have a layer of material (grass cuttings, plant leaves, vegetable tops, straw, etc.) 6 inches or more deep, scatter over it a coating of chemicals to aid decomposition (sulphate of ammonia, superphosphate, limestone) and water well. Forking over occasionally and wetting well during the season speeds decomposition. Place a 1-inch layer of earth at about 6-inch intervals to prevent bad odors. (Avoid using diseased plants or weeds, which should be burned.)

COSMOS
Semi-Double
Orange Ruffles



COSMOS
Yellow
Flare



PETUNIA
Scarlet



PETUNIA
Glamour



ZINNIA
Fantasy
Melody

GARDEN GUIDE POSTS




AQUILEGIA
(Columbine)



VERBENA
Brightness



DELPHINIUM
Wrexham
Giants



SCABIOSA
Tall Double
Mixed

(Below) NASTURTIUM, Golden Gleam



STORING VEGETABLES

Proper storage of root crops begins with planting, since they should just reach full maturity at the time frost is expected. (Maturity dates are given in each description.) Add 10 days to this for a margin of safety and then count back this period from the day killing frost can be expected. For example, Crosby's Egyptian beets mature in 60 days. If frost is expected Oct. 1st, then count back 70 days. This means that this variety should be planted for storage about July 20th.

In the case of cabbage, winter squash, pumpkins, onions and leeks, these should be planted 20 days ahead of the maturity dates given as they must be mature to keep well.

Never store bruised, rotting or imperfect vegetables. Quality decreases rather than improves with storage.

METHODS OF STORAGE

Beets, carrots, parsnips, salsify, turnips and rutabagas: Best stored in sand that is moist but not wet. Temperature as close to 33 degrees as possible, but must not go above 40. Growth begins at 42 degrees, which will destroy sweetness. Don't store all the parsnips indoors: leave some in the open garden for spring use. Freezing improves the flavor of these.

Cabbage, endive, celery and Chinese cabbage: Dig with earth clinging to roots, set on floor of cellar or in cold frame with earth packed around roots. Keep roots moist but never sprinkle tops. Allow slight circulation of air. Temperature 33 to 40 degrees.

Onions and potatoes: Dry between 33 and 40 degrees. Potatoes can stand a little more moisture than onions. After coming out of cold storage, potatoes will be better if kept in a room at 55 degrees for 10 days before using to get rid of sweet taste that develops at low temperature.


Pumpkins, Squash, Sweet Potatoes: Should be cured at 75 to 85 degrees for ten days, then kept in dry room at 55 degrees.



WHERE TO STORE

In House Cellar: Partition off corner of basement and control temperature by letting in cold air thru window. Cement floors will have to be sprinkled regularly or kept covered with damp sand. Insulate ceiling if room is under living space. Screen and ratproof to keep out pests.

Before vegetables are stored, wash floors and walls with household chlorine solution (Purex, Chlorox or Linc) to destroy spores that might rot vegetables.



Outdoors in Barrel: A barrel half sunk into the ground or dug into the side of a hill can be used to keep root crops in sand. Cover any exposed part of the barrel with two feet of straw in severe weather to keep out frost. Don't put all one kind of vegetable in the bottom, etc. Pack in mixed layers, so that an assortment can be taken out without emptying the whole barrel.

Cold Frame: A well-built cold-frame can be used to store many vegetables. Cabbage, endive, celery and Chinese cabbage can be planted in soil and covered with straw. Root crops can be piled in one end mixed with sand and covered. Put on sash and cover with mats. If glass is not covered, sun will keep frame too warm on bright days.

FOR BETTER FLOWERS FROM SEED

When and how to transplant: When the plantlets have fully developed their second pair of leaves, or "true" leaves as they are called, it is best to transplant them either to flats about three inches deep or to small pots. Remove them from the seed bed gently to damage the roots as little as possible and plant somewhat deeper than they originally grew, so that the stems will be well supported.

How to feed seedlings: When the seedlings have reached the stage of transplanting they require more plant food than they had in the seed flat mixture. Mix one teacupful of complete fertilizer with each bushel of soil mixture to make it fertile. One bushel of soil will take care of a great many plantlets.

Temperature and light: The transplanted seedlings should be kept at

a temperature of 45 to 50 degrees. If they are kept near a window, turn the boxes each day, otherwise the plants will lean towards the light and grow lopsided or spindly.

The right soil: Soil should be comparatively light in texture and preferably not too rich in plant food. One-third garden loam, one-third peat moss and one-third sharp sand will grow healthy seedlings.

When re-potting is necessary: As the little plants grow they may need to be shifted to larger pots to keep them in good growing condition up to the time the weather permits their being planted outdoors. This transplanting should be done when the roots have formed a mat around the inside of the pot. Usually it is best to change to a pot only one or two sizes larger than that in which the plant has been growing.

GARDEN SUPPLIES

"EROSIONET"

An inexpensive but strong, open-mesh fabric which is used in making new lawns where it is desirable to provide for the protection of young grass, or on steep banks where it will hold the soil and prevent washing and loss of turf by heavy rainfall. Also used for supporting Sweet Pea vines and shading cold-frames or perennial seed beds. 45 inches wide.

15c per linear yd., 100-yd. strips or more, 12c per linear yd.

FLOWER HOLDERS

Dome Type Nev-r-Tip. Heavy cast alloy, non-rusting. 1 1/4 in. diameter, \$0.25; 2 1/2 in. diameter, \$0.35; 3 3/8 in. diameter, \$0.55.

Pin Type Nev-r-Tip. Constructed with a heavy cast alloy base supporting sharp, needle-pointed, solid brass pins closely arranged. Round, 1 1/2 in. diameter, \$0.25; Round, 2 1/4 in. diameter, \$0.40; Round, 3 1/8 in. diameter, \$0.75.

GARDEN HOSE

VISIT OUR STORE for latest information and prices. We expect to be able to take care of your needs for garden hose, but cannot catalog definite qualities or prices because of the many uncertainties affecting this item.

GRAFTING WAX

Grafting Wax. The best on the market. 1/4-lb. pkg. 25c, 1/2-lb. pkg. 35c, lb.-pkg. 55c.

JARS (Preserving). Pint and quart sizes, self-sealing, high quality, for all types of home canning and preserving.

Cyclone Seed Sower—Uneven seeding is wasteful. The "Cyclone" will sow many acres per hour, any amount per acre. Does the work quickly and saves labor. Many hundred thousands of farmers in America, Canada and Europe have relied on "Cyclone" for sowing clover, timothy, alfalfa, lespedeza, sudan grass, oats, wheat, or any other seed that is sown broadcast. Each \$3.00.

Frost Can't Hurt Plants

Place HOTKAPS — patented paper hot-houses — over seeds or plants. Complete protection from destructive FROSTS, STORMS AND INSECTS. Ripen crops 3 weeks earlier; increase yield 18% to 51%. Maintain perfect mulch. Millions used by successful growers every season. Easy to set. 25 HOTKAPS for 50c. 100 for \$1.95. 250 for \$3.50. Setter free.



Germaco

HOTKAPS

A GERMACO PRODUCT

LABELS

Metal Plant Markers. Permanent metal labels, green enameled. 7 in. long, with white marking space 2 1/4 in. by 1 1/8 in. 12 for \$1.00; 25 for \$2.00; 100 for \$7.00.

Wood Pot Labels. The popular painted wood label for marking plants in pots or seed flats; 5/8-in. wide, 4-in. to 12-in.

	100	1000
4 in.	\$0.50	\$ 2.50
6 in.65	3.95
12 in.	1.75	13.95

100	1000	
6 in.65	3.95
12 in.	1.75	13.95

Wood Tree Labels, Wired. Painted wooden labels equipped with wires, for marking trees or shrubs; also used extensively for labeling Dahlia tubers.

100	1000	
3 1/2 x 5/8 in.	\$0.65	\$3.80

NOZZLES

Plastic Hose Nozzle. An adjustable, modernly designed red plastic nozzle. Light in weight, smooth finish, and adjustable to a complete variety of sprays and streams; also leak-proof shut-off. A very satisfactory substitute for the standard brass nozzle. Each 75c.

LAWN SPRINKLERS

We have Lawn Sprinklers which vary in price from \$1.00 up. We cannot however, tell just what styles or types we may have from day to day, throughout the season.

POTS AND ACCESSORIES

Standard Earthen Flower Pots. A wide range of sizes, from 2 1/2 in. to 12 in. diameter.

Earthen Flower Pot Saucers. 3 in. to 12 in.

Red Paper Pots. Made of water-proofed paper. Excellent for growing seedlings or for packing plants for shipment. Nicely finished and long lasting. 2 1/4 in. to 6 in. diameter.

Doz.	100	1000	
2 1/4 in.	\$0.20	\$0.85	\$ 5.25
2 1/2 in.23	1.00	5.75
3 in.28	1.25	6.85
4 in.40	1.65	9.45
5 in.45	2.35	13.75
6 in.50	2.60	17.00

SOIL TESTING

Sudbury Soil-Test Kits. Provide a simple method of determining the nitrogen, phosphorus and potash contents and also the degree of acidity. No knowledge of chemistry is needed. These kits are used by Government departments, commercial growers, and thousands of home gardeners.

Home Garden Kit (20 tests)	\$ 2.00
Club Model (50 tests)	4.75
Horticultural Size.....	9.75
Professional Size.....	18.50
Additional testing fluids: 2-oz. bottle, 75c; 6-oz. bottle, \$1.50.	

METAL PLANT PROPS



"Out since Pearl Harbor . . . now available again"

Green enameled steel props that provide perfect support for all types of flowers and plants.

Plant Props are rot-proof, termite-proof, wear-proof! They can be saved from year to year . . . look good in any part of the garden . . . and cost very little.

"America's Darling of the Props"

PLANT SUPPORTS

Bamboo Sticks, Green-Stained. These slim sticks are ideal for pot plants that require only light staking. Neat and inconspicuous.

	100	1000
18 in.	\$1.25	\$5.00
24 in.	1.40	7.00
30 in.	1.60	8.00
36 in.	1.75	9.50

Dahlia or Garden Stakes. Unpainted, square stakes made from Western Cedar.

Doz.	100	
4 ft., 3/4 in. square.....	\$1.95	\$13.50
5 ft., 3/4 in. square.....	2.50	16.50
6 ft., 3/4 in. square.....	2.95	20.50

Garden Stakes. Round stakes made from Western Cedar; stained dark green and pointed, the surface is ridged to prevent plant ties from slipping. Light weight, neat and almost rot-proof. 2 ft. to 8 ft.

TYING MATERIALS AND DEVICES

Raffia, Natural. A palm-fibre used for tying plants, bunching vegetables, weaving, and basketry. 1/2-lb. pkg. 50c. In braids (uneven weights) just as pulled from the bale, 95c per lb.; 10-lb. lots and over, 85c per lb.

Wayward Vine Guide and Support. An efficient and practical device for training Ivy or other vines on brick, stucco, stone or cement walls without defacing the wall. The support consists of a small tab with a copper wire hook for holding the vine. The tab is very easily cemented into place and will remain indefinitely; not affected by any weather conditions. Available in three colors to match any wall—Red for brick, Gray or White for stone or cement. Box of 25 (including cement) \$1.00.

WATERING DEVICES


Soil Soaker. Puts water just where water is needed—right in the soil at the plant roots. The Soil Soaker is made of porous canvas. Attached to a garden hose it fills up and, under normal pressure, water seeps through every pore of its entire length, soaking directly into the soil. 12 ft. long, \$1.40; 18 ft. long, \$1.90; 30 ft. long, \$2.90.

GARDEN TOOLS

At the time this catalog goes to press, there are still many uncertainties about the supply of garden tools. We have naturally done our best to gather an adequate supply of the tools you will need, and will be able to take care of most needs, but cannot promise complete assortments at all times.

(Prices are necessarily subject to change without notice)

CULTIVATORS



Speedy Cultivator. Highly tempered steel tines, sharp and curved to penetrate the soil with an easy pulling motion. Excellent for cultivating under shrubs. 4 tines, 4½ ft. handle, \$1.20



Victory Cultivator. Strong, rugged tool with forged steel teeth, adjustable fastening. Fine for plots not large enough to need a wheeled cultivator. 5 tooth, 4 ft. handle, \$1.50



Midget Cultivator Weeder. Short handle, 3 small prongs. Indispensable for close-in work.

Visit our store for further selection.

FORKS



Spading Fork. 4 angular back tines with polished face. Light weight. Bent handles, 11" tines, No. SD, \$1.60

Manure Fork. 4 polished oval tines 12½ in. long. Bent handle, No. A4, \$1.95



Hay Fork. 3 polished oval tines, 12 in. long. Bent handles, No. A34, \$1.60

Visit our store for further selection.



REVOLUTIONARY HAND SPRAYER

New HARCO is only professional-type Sprayer priced so low. Long extension rod sprays up, down, sideways—quickly penetrates all hard to reach places—without adjustment. Easy to use. Attractive, colorful. Only \$2.35.

HARCO SPRAYER
A Germaco Product

RAKES



Rakes, Curved Tooth, Level Head. 14 teeth on head 3½ x 14¾ in. Heads ¼ in. thick. 5½ ft. handles. No. S14, \$1.30
Wood Rakes for lawns. 28 in. wide, 14 teeth, 6½ foot handle, solid construction. Ask for prices.

Visit our store for further selection.

SCYTHES

Made of best crucible steel, strongly ribbed, finished in black with polished cutting edge.

Grass Scythe, for lawn work, \$2.25

Weed Scythe, strong and stiff, \$2.25

Bush Scythe, for underbrush, \$2.25

SHEARS

Grass Shears. Easy cutting action for trimming edges and borders. High quality steel, pinch-proof handles, easily operated. 12 in., \$1.00

Visit our store for further selection

SHOVELS and SPADES

Assortments and supplies are limited and we cannot tell what styles and sizes we will have from day to day. We suggest that you visit our store to make your selection.

TROWELS




All-Steel Garden Trowel. Made of one piece of rigid steel, with shaped, ventilated handle. Polished blade. \$1.00

Visit our store for further selection.

Weeders—Various types of approved weed cutters and pullers. See these handy labor-saving tools.

Turf-Edgers—Indispensable for keeping lawns neat and trim. Several styles.

LAWN MOWERS



Supplies are uncertain and we cannot tell what styles and sizes we will have from day to day during the season. We suggest that you consult us about mowers.

OTHER TOOLS

If you need other garden tools than those listed here, we suggest you consult us. Uncertainty and shortage of supplies makes cataloging of tools very difficult, but a visit to our store will place our services at your disposal.



Available supplies are limited. We will stock as full a line as may be obtainable. Visit our store.

INSECTICIDES AND FUNGICIDES

ANT TRAPS—For inside or outside of house. A safe container. (See Pest Exterminators.)

ANT BAIT—A bait that needs no feeders or traps. Easy to use in handy tubes. 1-oz. 35c; 3-oz. 85c. (See Pest Exterminators.)

ARSENATE OF LEAD. The standard insecticide for controlling practically all leaf-eating or chewing insects. Can be used as a spray or dust. 1 to 2 tablespoonfuls makes 1 gallon spray.

BLACK LEAF 40. Nicotine Sulphate 40% solution. For controlling aphis (plant lice), thrip and other soft bodied sucking insects. 1 teaspoonful makes 1 gallon. 1 oz. bottle 36c; 5 oz. \$1.05.



ACME ROTENONE GARDEN GUARD is an ideal insecticide for the home garden. Dust or spray. The killing ingredient is Rotenone — non-injurious to human and warm blooded animals. Acts both as contact and stomach poison against a wide range of insects. 1 lb. sifter carton, 35c. 1 lb. pump gun. 60c.

ACME RED RIVER POTATO MIX—dust or spray. A scientific formula combining a patented copper fungicide with quick-killing arsenic. Kills potato bugs quicker—drives off flea beetle and leaf hopper—prevents blight—stimulates foliage. Get top prices for your harvest.

ROTELONE GARDEN SPRAY (NNOR)—see Sucking Insects.

ACME TOMATO DUST—see Blight and Fungus.

ACME VEGETABLE AND FLOWER GARDEN DUST—see Sucking Insects.

ACME APHIS SPRAY—see Sucking Insects.

PARIS GREEN—see above.

ARSENATE OF LEAD—see above.



ROTELONE GARDEN SPRAY (NNOR) containing Rotenone kills most common garden bugs of both sucking and leaf eating types. You've got to wet them — to get them. NNOR penetrates the waxy or hairy covering, brings the tremendous power of Rotenone into action. 1 oz., 35c; 6 oz., \$1.00; 16 oz., \$2.15; Gals. \$13.50.

ACME EMO-NIK is a valuable spray for the control of scale, red spider, mealy bug, white flies, rust mites, aphis, and many other common insects. A complete contact insect spray combining nicotine in its most active form with Emo, a high quality summer oil emulsion. Practically odorless when

BORDEAUX MIXTURE. Standard copper fungicide for controlling black spot, mildew, blight, leaf-curl, scab and other fungous diseases on flowers, fruits, vegetables and other plants. 1 lb. makes 5 gallons. 1 lb. 35c; 4 lbs. 95c.

CALCIUM ARSENATE—Arsenate of Lead and Lime. A cheap quick-acting poison for potato bugs and other leaf-eating insects.

CERESAN — Effective in controlling covered smut and stripe of barley, oat smuts, and bunt or stinking smut of wheat. A pound treats 32 bushels. 4-oz. pkg. 30c; lb. 80c; 4-lb. pkg. \$2.70; 25-lb. pkg. \$15.00.

sprayed. 4½ oz., 35c. Pint, 65c. Quart, 95c. Gallon, \$2.50.

ACME APHIS SPRAY combines a soft spray soap with Black Leaf 40. Recommended for aphis (plant lice), thrip, leaf hopper, and many more hardy insects infesting rose bushes, flowers, shrubs, vines, bushes, and trees. 3 oz. collapsible tube 35c. 12 oz., \$1.00. 2½ lbs. \$2.35.



ACME VEGETABLE AND FLOWER GARDEN DUST. A dust or spray containing nicotine for many common varieties of sucking insects, and arsenicals for leaf eating insects for the home vegetable and flower garden. 1 lb. cartons, 40c. 4 lb. bags, 1.10.

ACME ROTENONE GARDEN GUARD—see Chewing Insects.

LIME SULPHUR—see above.



ACME TOMATO DUST is especially prepared to control blight and worms preying on tomatoes. While specifically designed for tomatoes, it also is equally effective in the control of leaf chewing insects and blight on many other vegetables such as potatoes, cucumbers, beans and cabbage, also many flowers as roses, azalea, evergreens, asters, and hollyhock. 1 lb. sifter carton, 35c. 1 lb. pump gun, 55c. 4 lb., 80c. 50 lb., \$9.00.

ACME KOPPER QUEEN (Mildew Spray) is a stainless liquid copper spray for the control of mildew, leaf spot, and black spot on rose bushes, flowers, and many other kinds of foliage. Leaves no unsightly residue. A liquid spray replacing Bordeaux Mixture and Sulphur. ½ pt., 35c; Pint, 60c; Qt. 90c; Gal., \$2.25.

ACME WETTABLE DUSTING SULPHUR is an improved dusting sulphur for home use which can also be used as a liquid spray. Controls red spiders on evergreen, mildew, leaf spot, black spot, and rust on rose bushes, chrysanthemums, snapdragons, carnations, and certain other foliage. 98% passes through a 325 mesh screen. An excellent chigger chaser.

ACME RED RIVER POTATO MIX—see Chewing Insects.

BORDEAUX MIXTURE—see above.

CYANOGAS "A" DUST. Calcium cyanide for killing rats, moles, gophers, etc. Gives off deadly gas but leaves no poisonous residue. ¼ lb. can 25c; ½ lb. 45c; 1 lb. 75c; 5 lbs. \$3.00; 25 lbs. \$10.00.

CYANOGAS ANT KILLER—A. Dust —A dry powder to be blown into ant-hills, etc. 4 oz. can 30c.

CYANOGAS G. Fumigant—The standard for greenhouses, mushroom houses, and for soil-fumigation. 5 lb. can \$3.00; 25 lb. can \$10.00.

COPPER SULPHATE (or Blue Stone)—Fine granulated.

DORMANT OIL SPRAY—A self-emulsifying spray-oil that kills scale aphid, psylla, red mite, leaf-hopper, and certain other insects that overlive Winter on trees. The only effective control for scale insects. Will not injure paint.

DOG-CHECK—An effective proven curb to canine habits, needed by all whose neighbor's dogs "visit" and wither flowers, shrubs and saplings. One spraying usually lasts from two to three weeks — unaffected by light rains. Each package contains 6 DOG-CHECK APPLICATORS. 60c and \$1.00 retail sizes.

FISH-OIL SOAP—An old fashioned remedy for sucking and scale insects.

MOLE KILLER—See Pest Exterminators.

MOUSE POISONS—See Pest Exterminators.

NAPHTHALENE FLAKES—Frequently mixed with soil in greenhouses and gardens to destroy slugs and wire-worms. Also useful in controlling thrips on gladiolus bulbs.

NICO-FUME LIQUID—A greenhouse fumigant and spray material. 1 lb. can \$2.50; 4 lb. can \$7.10; 8 lb. can \$13.00.

NICO-FUME POWDER—A greenhouse fumigant for the control of numerous insect pests. ½ lb. and 1 lb. size packed in pressure fumigator containers. ½ lb. tins 60c; 1 lb. tins \$1.00; doz. \$9.00.

RAT POISONS—See Pest Exterminators.

FLEA-NOT

Kills fleas, lice and other parasites that infest the skins of dogs and other pets. Absolutely harmless to the skin of animals. Easy to use. Put up in an attractive shaker top can. 25c Retail Size.

Acme Scientific Rose and Flower Spray



A "system" of spray treatment for roses and flowers of all kinds. Eliminates guess work. Each set contains three separate items — each scientifically made to combat one of the three major classifications of blight and insect pest. Mix all three together with water. Use regularly. Result: complete all-around protection. Surprisingly economical.

Sizes	Amount of Spray	No. of Roses Protected all season	Prices
No. 3. .12 qts...	7 to 12 bushes	.00	\$0.95
No. 6. .24 qts...	15 to 25 bushes	.00	1.50
No. 24. .96 qts...	60 to 100 bushes	.00	4.00
No. 48. .48 gals...	120 to 200 bushes	.00	6.00
No. 96. .96 gals...	240 to 400 bushes	.00	9.75

PARIS GREEN. An effective insecticide for control of chewing insects, particularly recommended for potato bugs. $\frac{1}{4}$ lb. 20c; lb. 55c.

POTATO MIX — See "Chewing Insects."

RODENT KILLER—Molite — The new gas type cartridge for killing moles, rats, gophers and other underground rodents. Simply light fuse and place in runways or tunnels. The poison gas brings quick death to rodents. Non-explosive, non-inflammable. Safe to use. 15c each; \$1.50 doz.

ROtenone DUST. The greatest advancement in years in the insecticide field. Rotenone is a non-poisonous dust that effectively controls nearly every kind of insect, whether chewing or sucking. Particularly recommended for bean beetles, cabbage worms, aphids. Offered only for vegetables in accordance with W.P.B. ruling.

RED ARROW GARDEN SPRAY — A highly concentrated Pyrethrum and Rotenone extract containing soap, spreader and sticker. Harmless to animals. 1 oz. 35c; 4 oz. \$1.00; 1 pint \$2.85; 1 quart \$5.00; 1 gallon \$17.00.

SEMESAN — Effective in controlling damping-off, scab, hard rot, etc. 2 oz. can 40c; 1 lb. can \$2.50; 5 lb. can \$11.40; 25 lb. can \$52.50.

SEMESAN BEL — For use on potatoes. Controls such seed-borne disease organisms as scab, rhizoctonia, and black-

leg. One pound treats 16 to 20 bushels of seed. 2 oz. 30c; lb. \$1.65; 5 lbs. \$7.15; 25 lbs. \$33.25.

SEMESAN JR. (New Improved). Disinfects corn against ear rot and stalk diseases. 2 oz. treats 1 bushel seed corn. $1\frac{1}{2}$ oz. 15c; 1 lb. 75c; 5 lbs. \$3.00.

SLUG SHOT. Famous non-poisonous dust for killing potato bugs, cabbage worms and lice and other insects and worms on vegetables. 1 lb. 30c; 5 lbs. \$1.25; 25 lbs. \$5.50.

SULPHUR, DUSTING — For controlling powdery mildew, blight, black spot, rust and most rose diseases.

SULPHUR, WETTABLE — An improved sulphur which can also be used as a liquid spray. 98% passes through 325 mesh screen.

TOBACCO DUST, Gold Leaf — Finely pulverized. Used for dusting plants to repel various insects. Guaranteed to contain not less than 1% nicotine by weight, dry basis.

TREE TANGLEFOOT — Perfect safeguard for trees, vines, and shrubs against gypsy, brown-tail and tussock moths, canker worms, ants, and other creeping insects. 6 oz. can 45c; 1 lb. can 80c; 5 lb. can \$3.75; 10 lb. can \$7.00; 25 lb. can \$15.00.

VOLCK, Nursery — An insecticide effective against mites, scale, red spider and mealy bug, and when combined with "Black Leaf 40," Volck also controls thrips, lacy-wing fly and chrysanthemum midges. $\frac{1}{2}$ pt. can 35c; pt. can 50c; qt. can 75c.

TRI-OGEN FOR ROSES

TRI-OGEN SPRAY — three way treatment for roses and most vegetable plants. Used and highly recommended by leading rosarians. Controls insects and fungus, assuring healthy, strong and vigorously blooming roses. **E Kit** 90c; **A Kit** \$1.50; **B Kit** \$4.00; **C Kit** \$6.00; **D-Estate kit** \$20.00

TRI-OGEN DUST — a perfected dust for three-way control of insects and fungus diseases, for those who prefer to dust. Especially good for roses but useful on all flowering and vegetable plants. **1 lb. 65c; 5 lbs. \$3.00**



Spray with BLACK LEAF 40

Aphis (plant lice), leafhoppers, leaf miners, mealy bugs, lace bugs, most thrips, young sucking bugs and similar insects can be controlled.

An ounce makes 6 gallons of spray effective on aphis. Full directions with package — A quick, sure, economical control for these insect pests.

Just a little spread on roosts kills poultry lice and feather mites.


ZOTOX CRAB GRASS KILLER

Destroys Crab Grass in lawns without permanent injury to lawn grasses. Also kills Chickweed, Spurge, Dandelion, Plantain and many other lawn weeds. A 16 oz. bottle makes 10 gallons and treats up to 2000 sq. ft. Price: 8 oz. bottle \$1.00; 16 oz. \$1.50; 32 oz. \$2.50; 5 lb. \$4.50.

WEEDONE
TRADE MARK

The new scientific weed killer. Kills Poison Ivy, Poison Oak, Bindweed, Jap. Honeysuckle, other noxious weeds. Sprayed on leaves it travels down and kills right out to the root tips. Non-irritating, non-corrosive, contains no arsenic or chlorates. Just mix in water and spray.

6-oz. bottle (covers 300 sq. ft.)	\$1.00 — 1-qt. bottle (covers 1500 sq. ft.)	\$4.00 — 1-gal. bottle (covers 6000 sq. ft.)	\$12.00 —
5-gal. can (covers 30,000 sq. ft.)	\$55.00 .		



SPRAYERS and DUSTERS




The supply of garden sprayers is still critical, but we hope to be able to take care of your needs.

Good spraying equipment is vital to the success of the garden, so we are making every effort to maintain our supplies for you. Consult us on all types of spray equipment.

Compressed Air Tank Sprayers

Open top or funnel top, 2 to 4 gallon sizes, for all types of farm and garden spray work, whitewashing, etc. Easy to fill, easy to clean.

Continuous Sprayers




Convenient, efficient sprayers for insecticides, fly oils, deodorants, fly repellents, stock sprays, etc. For farmer, gardener, poultryman.

Light Hand Sprayers

The right models for the back yard garden, chicken house, etc. (Supply is short, but we can probably take care of you.)

Dusters



Various models, made to handle any powdered insecticide, with any desired discharge from a very fine to a very heavy cloud. Heavy duty models for truck farms, orchards, etc. Lighter models for the garden.

Wheelbarrow Type Sprayers



Heavy duty portable outfits for orchard, farm, and industrial work. Consult us about models available.

Bucket Spray Pumps

Bucket or barrel types for whitewash, cold water paints, insecticides, caustic mixtures, etc. Consult us about models available.

PLANT FOODS

ROOTONE

Reg. U. S. Pat. Off.

The Plant Hormone Powder

This nationally known hormone powder makes cuttings root faster and more successfully. It stimulates seed germination so that rooting is heavier and you get a better stand of seedlings. Endorsed and used by leading growers everywhere. Trial packet 25c—2 oz. jar \$1.00—1 lb. can \$5.00.

TRANSPLANTONE

Reduces shock and wilt when plants are moved. Produces earlier maturity on tomatoes, lettuce, celery, peppers, tobacco and cabbage. Used on plants, shrubs and trees it encourages growth and bloom. One teaspoonful of powder makes 10 gallons of solution. Trial packet 25c—1 oz. can 50c—3 oz. can \$1.00—1 lb. can \$4.00.



MEET THE NEW, IMPROVED PLANT FOOD WONDER!

The same inexpensive Plantabbs with full pre-war quota of Nitrogen

You can't beat Plantabbs for growing bigger, better garden flowers, potted plants and vegetables. They're ready for you now in new, WHITE tablet form. Each tablet contains a full, rich meal of Nitrogen, Phosphoric Acid, Potash (11.15.20) plus Vitamin B₁. They're clean, odorless, safe and easy-to-use. Just push a tablet into the soil alongside each plant every 10 days. The usual watering or rain will quickly dissolve it. If you prefer liquid, simply drop 4 tablets into 1 gallon of water, and there you have it. No matter how you use Plantabbs, you'll be amazed at the difference they make in your plants. Start feeding your potted plants, garden flowers and vegetables by this modern, convenient method. If you don't get wonderful results, you'll get your money back. The new, improved Plantabbs come in four economical sizes — 25c, 50c, \$1, \$3.50.

FULTON'S
PLANTABBS
Odorless Plant Food Tablets

While there are still some wartime restrictions on the manufacture and sale of fertilizers, the situation is greatly improved. This spring there should be plenty of suitable fertilizer for every home garden.

Ask us about it!

Whatever your needs for plant foods may be, consult us about them. We'll see to it, as usual, that your garden and lawn get the food-help they need.

ALUMINUM SULPHATE — A material used in creating acid soils necessary for successful culture of Azaleas, Rhododendrons, and similar plants.

ASHES, HARDWOOD — High potash content. Contains all of the fertilizer elements except nitrogen. Tends to drive away insects.

BACTO — Converts leaves, grass clippings, table scraps, etc., into rich fertilizer-humus in a few weeks. 5-lb. treats half a ton.

BONE MEAL — All-purpose fertilizer, rich in the two principal plant foods—nitrogen and phosphoric acid.

FRUITONE — The remarkable hormone spray that stops pre-harvest drop of fruit. Sprayed on flowers of tomatoes it makes a high percentage of the crop **SEEDLESS!** Also beneficial on beans and on flowers.

G & O ROSE FOOD



... at last, a real organic, no-filler fertilizer, **ESPECIALLY FOR ROSES!**

You can't feed roses a "good-for-everything" commercial food and get the color, size of bloom and stock you want!

G & O Rose Food is specialized — a balanced combination of ORGANIC ingredients — selected for the particular "make-up" and sequence availability that roses need. TRY IT—and see what a difference it makes!

For better color, size, strength

Stretches Fertilizer!

Amazing! Bacto will multiply 1 lb. of chemical fertilizer into 60 lbs. of better, non-burning organic-type fertilizer. Easy! Quick!

Hastens Compost

Don't burn leaves! Bacto makes rich fertilizer-humus from leaves, table scraps, grass clippings—even sawdust—in a few weeks. 5-lb. treats $\frac{1}{2}$ -ton.

BACTO also activates hot beds and pot soils; reduces odor in outdoor toilets—a bacterial product of many valuable uses. (Made by Jean Maclean, Bridgeton, Ind.)

5-lb. 169 25-lb. 549
Ship. wt. 6 lbs. Ship. wt. 27 lbs.
Trial can, 25c 3 for 71c



HUMUS — One of the best soil conditioners. Makes soil more workable and helps it hold moisture. Not a substitute for fertilizer, but an excellent companion for it. In convenient sized packages.

LIME (Hydrated) — Sweetens sour soil for lawns, and clover. Is used as a reducer and in combination of many insecticides; for whitewash, deodorizing.

MURIATE OF POTASH — Improved Potash. Used liberally for all root crops.

NITRATE OF SODA — Very quick source of nitrogen. Hastens maturity of crops. Should not be applied until plants are well above ground.

PLANTABBS — Clean white odorless plant tablets for indoor or outdoor use, combine high analysis food value with a plentiful supply of Vitamin B-1.

PEAT MOSS — Fine for garden mulch, as it conserves moisture. When incorporated in the soil, it promotes growth and supplies humus, which is very essential for growing plants. Fine winter protection for lawns; also as a mulch on strawberry perennial beds, roses, evergreens, etc.

SHEEP MANURE — An economical and effective balanced plant food. Excellent for lawns, gardens and general use.

SOIL TEST KITS

Do your own soil testing! Kits are simple, practical, easy to use. Let you determine just how to fertilize and lime for best results. Complete kit, good for 20 individual tests, \$2.00.

SULPHATE OF POTASH — Highly recommended for plants requiring acid soil.

SUPER PHOSPHATE — Aids plant nutrition and assists the plant to assimilate other ingredients. An efficient fertilizer element for lettuce, truck crops, cereal crops and alfalfa.

TOBACCO STEMS — Used as a mulch for roses, asters, lettuce, etc., and for fumigating plants to kill insects.



ROOTONE—You can help cuttings to root faster by dipping them in the hormone powder, Rootone, before planting. Percentage of successful rooting is increased remarkably by this new treatment.

TRANSPLANTONE — Remarkable new Vitamin-Hormone stimulant effective in reducing transplant shock, and wilt of all kinds of plants or shrubs. Also effective in bringing about earlier maturity on tomatoes, lettuce, celery, peppers, tobacco and cabbage.

LEGUME INOCULATION — Just as necessary for garden peas and beans, lima beans, sweet peas and lupines as for alfalfa, clover, etc. The nitrogen-gathering bacteria build up the soil, increase your yields. Inoculation is the cheapest form of crop insurance. Packages from the 10c garden size up to 5-bushel size.

FEEDING THE GARDEN

• How much fertilizer? In general, the use of from 25 lbs. to 35 lbs. of a good general fertilizer like 4-12-4 or 5-10-5 to every thousand square feet of garden space will take care of all the needs of most plants.

• The addition of compost will make soil easier to work and add certain minor elements not available in commercial plant foods.

• Sheep or other animal manures are particularly valuable in sandy soils because they add soil bacteria usually missing in such soils.

• In growing root crops and bulbs the use of 4 to 5 lbs. of muriate or sulphate of potash to every 1,000 square feet will make for firmer bulbs, tubers and roots. Wood ashes can be substituted for these (except on potatoes, where the higher lime content of wood ashes encourages scab), applying 12 lbs. per 1,000 square feet.

• When seeds are planted at temperatures below 42 degrees (i.e. lettuce, spinach and grass seed sown in early spring) the treatment of the seed with ROOTONE supplies hormones that substitute for the natural plant hormones which are not active at low temperatures.

Notes on INSECTS AND DISEASES

It is not possible to give complete directions for spraying for insects and diseases, but if you can bring in a specimen showing the difficulty we shall be glad to do all we can to recommend a suitable spray or dust.

• Chewing insects cut or tear out pieces of leaves or stems. They are killed by stomach poisons like calcium arsenate, arsenate of lead or Paris Green. When the insects are on plants that are to be eaten, Rotenone or Pyrethrum are safer, but do not always give as complete control, as the insects themselves must be hit directly.

• Sucking insects stab the stem or leaf and suck out the plant juices and cannot be poisoned through their stomachs. They are smothered by sprays or dusts that paralyze their breathing apparatus. Rotenone, Pyrethrum and Nicotine Sulphate are commonly used for this purpose.

• Fungicides control most plant diseases, which are caused by fungi or moulds. Bordeaux mixture, dusting sulphur and lime sulphur are used for this purpose, while for seeds and potato tubers, organic mercury compounds like Semesan give good results.

HYPODEX PLANT FOOD

Grows bigger and better house plants, flowers, vegetables, lawns and trees in poorest soil—even in sand, cinders, or water. Results are sure—it doesn't burn.

Clean, odorless, soluble powder—complete and balanced.



1 oz. package 10c
(makes 6 gallons)
3 oz. package 25c
1 lb. package \$1.00
(makes 100 gallons)

Available in drums, write for prices.

• Compound spray materials are valuable in the smaller garden, since they combine fungicides and insecticides, or materials for controlling both chewing and sucking insects.

• **CAUTION:** Always use insecticides as directed on the package: never stronger than recommended. Avoid using materials containing sulphur when the sun is shining brightly and when the air temperature is above 80 degrees, as they will burn badly at temperatures above this point.

WHEN IN DOUBT, ASK US! Whatever your questions about insect pests or plant feeding—don't hesitate to consult us!

A Dry Bath for your dog



Avoid the fuss and muss of bathing your dog by cleaning quickly, efficiently and thoroughly with Sudbury Hygienic Cleaning powder. Recommended by authorities . . . helps rid fleas and mites . . . leaves coat soft and silky. Giant size, \$1.00.

Liquid Chaperone, \$1.00—Keeps dogs away from flowers, shrubs.

Powder Chaperone, \$1.00—Keeps dogs off furniture.

DOES YOUR DOG Itch.. Scratch?



Don't Blame Fleas, Mange or Diet . . . 2 to 1 It's "Fungitch"
Amazing New Sulfadene Eliminates Canine Skin Infection Long a Puzzle to Science

IF YOUR dog or cat scratches, it probably has "FUNGITCH" . . . a fungus infection caused by contact with mouldy grass and weeds. "FUNGITCH" starts as an itch . . . then leads to dandruff-like scales or sores with brownish edges, mouldy odor—or infection in ears, on paws, underbody, tail or eyes . . . resulting in something more serious, expensive care, or actual loss. Treatment of 3,121 dogs and cats—at Dr. Merrick's Animal Hospital—reveals SULFADENE stops itching quickly, clears up sores in a few days. To relieve your pet apply SULFADENE externally at first sign of "FUNGITCH." Only 98c for generous treatment. Guaranteed.

Dr. MERRICK'S SULFADENE

PEST EXTERMINATORS... "Their Last Meal"

ANT-X JELLY BAIT—A most effective bait for destroying all species of ants—SIMPLE TO USE—Not a repellent but a food bait that ants eat and carry back to the nest. Destroys entire colony. "THEIR LAST MEAL." 35c and \$1.00 retail sizes.

ANT-X ANT TRAPS—A safe, sanitary method for controlling ants in house and garden. Attracts and destroys both sweet and grease-eating ants. Cannot be opened by children or spilled by pets. ANT-X TRAPS may be placed inside ice-boxes or other food containers without contaminating foods. "THEIR LAST MEAL." 25c retail size.

RAT-NOTS—A non-poisonous prepared bait containing genuine RED SQUILL. A safe, sanitary way to exterminate rats. Rats are attracted to RAT-NOTS because of their tempting odor. "THEIR LAST MEAL." 25c and \$1.00 retail sizes.

MOUSE-NOTS—A meaty, easily hulled, treated seed bait which appeals to the finicky taste of mice. Simply spread in places where mice frequent. "THEIR LAST MEAL." 25c and \$1.00 retail sizes.

ROACH-NOTS—A prepared powder bait, quick and effective. ROACH-NOTS "Lays them on their backs." Also quickly kills waterbugs and silverfish. "THEIR LAST MEAL." 25c and \$1.00 retail sizes.

MOLE-NOTS—Life-saver for lawns, flower and seed beds. Sure death to bulb-eating, root-destroying moles. MOLE-NOTS safeguard bulbs when placed in ground at time of planting. "THEIR LAST MEAL." 35c, \$1.00 and \$2.00 retail sizes.

RAT-NOT-PASTE—Recommended for big, well-fed, hard-to-catch rats which dodge cats and traps,

and are finicky in their eating habits due to easy access to tasty-garbage, and feed and food warehouses. Ideal, too, for control of chipmunks, squirrels, prairie dogs, etc. "THEIR LAST MEAL." Packaged in handy, convenient tubes. 35c and \$1.00 retail sizes.

SKEET-CHECK—An ointment repellent, packaged in convenient handy tubes, easy to apply and pleasant to use. Unexcelled for repelling mosquitoes, gnats, black flies, deer flies and other annoying insect pests. SKEET-CHECK is also an excellent soothing salve that will help to allay swelling when bitten. 35c retail size.

KILMICE—A treated, tasty, impregnated seed bait which quickly destroys mice in house and garden. Easy to use; clean, sanitary, no mixing—no traps needed. "THEIR LAST MEAL." Large 2½ oz. package 25c.


Complete Garden Service—AT OUR STORE OR BY MAIL

EVERETT SEED CO.

78 Alabama Street, S.W. (Between Whitehall and Pryor)

ATLANTA, GA.

**You KNOW you're safe...
with EVERETT Farm Seeds**



**Did you ever stop to think—"just how does a seedsman
build a reputation for himself?"**

**One thing is sure—fancy selling talk can't do the job!
At least, not for long!**

**No sir, there's just one way to make a lot of people say,
year after year, "there's a seedsman you can depend on!"
And that one way is to **SELL SEED THAT'S RIGHT!****

**People feel that way about Everett's seeds. The longer
they buy here, the more they know that we are **ALWAYS**
careful and conscientious!**

**We value that reputation above all else—and you can
be mighty sure we'll never knowingly do anything to
hurt that reputation.**

**IF YOU HAVEN'T ALREADY FOUND THAT OUT FOR
YOURSELF, THIS IS A GOOD TIME TO START!**

**Cow Peas
Soy Beans
Peanuts
Velvet Beans
Crowder Peas
Lespedeza
Clover
Pasture Grasses
Sorghum Cane
Sudan
Millet
Field Corn
Alfalfa
Hegari
Kaffir
Milo
Crotalaria
Austrian Peas
Vetch
Lupine
Oats
Rye
Barley
Wheat**